Please	type a	plus	sian	(+)	inside	this	box	_
	.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	piuc	U.G.,	١.,	Horac	uno		

PTO/SB/05 (12	/97) (modif
Approved for use through 09/30/00. O	MB 0651-0
Patent and Trademark Office: U.S. DEPARTMENT OF	COMMER

i	
	UTILITY
	PATENT APPLICATION
	TRANSMITTAL

Attorney Docket No.	348022001200	Total Pages	57
Firs	t Named Inventor or Application i	ldentifier	<b>N</b>
e Chao Yu and Danie	al P. Henderson		Foot

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Express Mail Label No. EL 034 669 514 US

#### CERTIFICATE OF MAILING BY "EXPRESS MAIL"

Express Mail Label No.: EL 034 669 514 US

Date of Deposit: September 9, 1999

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is april commissioner for Patents, Washington, D.C. 20231.

Assistant Commissioner for Patents **APPLICATION ELEMENTS** ADDRESS TO: **Box Patent Application** See MPEP chapter 600 concerning utility patent application contents. Washington, DC 20231 Fee Transmittal Form 1. 🔀 Microfiche Computer Program (Appendix) (Submit an original, and a duplicate for fee processing) Specification [Total Pages 53 ] Nucleotide and/or Amino Acid Sequence Submission 2. 🗴 (preferred arrangement set forth below) (if applicable, all necessary) - Descriptive title of the Invention - Cross References to Related Applications b. Paper Copy (identical to computer copy) - Statement Regarding Fed sponsored R & D c. Statement verifying identity of above copies - Reference to Microfiche Appendix - Background of the Invention **ACCOMPANYING APPLICATION PARTS** - Brief Summary of the Invention - Brief Description of the Drawings (if filed) 8. Assignment Papers (cover sheet & document(s)) - Detailed Description Power of Attorney 37 CFR 3.73(b) Statement - Claim(s) (when there is an assignee) - Abstract of the Disclosure 10. 3. 🔀 English Translation Document (if applicable) Drawing(s) (35 USC 113) [Total Sheets 27] Information Disclosure 4. 🔲 Copies of IDS Oath or Declaration [Total Pages Statement (IDS)/PTO-1449 Citations a. U Newly executed (original or copy) 12.L **Preliminary Amendment** Return Receipt Postcard (MPEP 503) b. Copy from a prior application (37 CFR 1.63(d) (Should be specifically itemized) (for continuation/divisional with Box 17 completed) [Note Box 5 below] 14. Small Entity Statement filed in prior application, LI DELETION OF INVENTOR(S) Statement(s) Status still proper and desired Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b) 15. Certified Copy of Priority Document(s) (if foreign priority is claimed) Incorporation By Reference (useable if Box 4b is checked) The entire disclosure of the prior application, from which a copy of the 16. oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein. 17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information: ☐ Continuation ☐ Divisional Continuation-in-part (CIP) of prior application No: 18. CORRESPONDENCE ADDRESS Catherine M. Polizzi Registration No. 40,130 Morrison & Foerster LLP

J ū î. ũ r. fl Į

755 Page Mill Road

Palo Alto, California 94304-1018 Telephone: (650) 813-5651 Facsimile: (650) 494-0792

If a paper is untimely filed in the above-referenced application by applicant or his/her representative, the Assistant Commissioner is hereby petitioned under 37 C.F.R. § 1.136(a) for the minimum extension of time required to make said paper timely. In the event a petition for extension of time is made under the provisions of this paragraph, the Assistant Commissioner is hereby requested to charge any fee required under 37 C.F.R. § 1.17(a)-(d) to **Deposit Account**No. 03-1952. However, the Assistant Commissioner is NOT authorized to charge the cost of the issue fee to the Deposit Account.

The filing fee has been calculated as follows:

FOR	NUMBER FILED	NUMBER EXTRA	RATE	CALCULATIONS
TOTAL CLAIMS	28 - 20 =	8	x \$18.00	\$224.00
INDEPENDENT CLAIMS	2 - 3 =	0	x \$78.00	\$0.00
MULTIPLE DEPENDENT	\$0.00			
	enski prograf Libria (2005) Standardski sekologija Postaj postaj sekologija	<del>of Bos (All money prop</del> er on the bea	BASIC FEE	\$760.00
TOTAL OF ABOVE CALCULATIONS =				\$984.00
Reduction by 1/2 for filing by small entity (Note 37 C.F.R. §§ 1.9, 1.27, 1.28). If applicable, verified statement must be attached.				\$492.00
The same of the sa	RARANIANA Managaran Jawa Sanasa Managaran Managaran	je i se konstantije in die Georgiaa konstantij g	TOTAL =	\$492.00

Applicant(s) hereby petitions for any required relief including extensions of time and authorizes the Assistant Commissioner to charge the cost of such petitions and/or other fees or to credit any overpayment to **Deposit Account No. 03-1952** referencing docket no. <u>348022001200</u>. A duplicate copy of this transmittal is enclosed, for that purpose.

Dated: September 9, 1999

Respectfully submitted,

Catherine M. Polizzi Registration No. 40,130

Morrison & Foerster LLP 755 Page Mill Road

Palo Alto, California 94304-1018

Telephone: (650) 813-5651 Facsimile: (650) 494-0792

#### CERTIFICATE OF MAILING BY "EXPRESS MAIL"

Express Mail Label No.: EL 034 669 514 US

Date of Deposit: September 9, 1999

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231.

Shirley Labor

## ADENOVIRUS VECTORS CONTAINING CELL STATUS-SPECIFIC RESPONSE ELEMENTS AND METHODS OF USE THEREOF

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the priority benefit of U.S. Provisional Patent Application No.60/099,791, filed September 10, 1998. The priority application is hereby incorporated herein by reference in its entirety.

# STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

(Not Applicable)

20

15

5

10

#### TECHNICAL FIELD

This invention relates to cell transfection using adenoviral vectors. More specifically, it relates to cell status-specific replication of adenovirus vectors in cells, regardless of tissue or cell type.

25

#### **BACKGROUND ART**

In spite of numerous advances in medical research, cancer remains the second leading cause of death in the United States. In the industrialized nations, roughly one in five persons will die of cancer. Traditional modes of clinical care, such as surgical resection, radiotherapy and chemotherapy, have a significant failure rate, especially

for solid tumors. Neoplasia resulting in benign tumors can usually be completely cured by removing the mass surgically. If a tumor becomes malignant, as manifested by invasion of surrounding tissue, it becomes much more difficult to eradicate. Once a malignant tumor metastasizes, it is much less likely to be eradicated.

5

Excluding basal cell carcinoma, there are over one million new cases of cancer per year in the United States alone, and cancer accounts for over one half million deaths per year in this country. In the world as a whole, the five most common cancers are those of lung, stomach, breast, colon/rectum, and uterine cervix, and the total number of new cases per year is over 6 million.

10

Lung cancer is one of the most refractory of solid tumors because inoperable cases are up to 60% and the 5-year survival is only 13%. In particular, adenocarcinomas, which comprise about one-half of the total lung cancer cases, are mostly chemo-radioresistant. Colorectal cancer is the third most common cancer and the second leading cause of cancer deaths. Pancreatic cancer is virtually always fatal. Thus, current treatment prospects for many patients with these carcinomas are unsatisfactory, and the prognosis is poor.

15

Hepatocellular carcinoma (HCC or malignant hepatoma) is one of the most common cancers in the world, and is especially problematic in Asia. Treatment prospects for patients with hepatocellular carcinoma are dim. Even with improvements in therapy and availability of liver transplant, only a minority of patients are cured by removal of the tumor either by resection or transplantation. For the majority of patients, the current treatments remain unsatisfactory, and the prognosis is poor.

20

Breast cancer is one of the most common cancers in the United States, with an annual incidence of about 182,000 new cases and nearly 50,000 deaths. In the industrial nations, approximately one in eight women can expect to develop breast cancer. The mortality rate for breast cancer has remained unchanged since 1930. It has increased an average of 0.2% per year, but decreased in women under 65 years of

age by an average of 0.3% per year. See e.g., Marchant (1994) Contemporary Management of Breast Disease II: Breast Cancer, in: Obstetrics and Gynecology Clinics of North America 21:555-560; and Colditz (1993) Cancer Suppl. 71:1480-1489.

5

Despite ongoing improvement in the understanding of the disease, breast cancer has remained resistant to medical intervention. Most clinical initiatives are focused on early diagnosis, followed by conventional forms of intervention, particularly surgery and chemotherapy. Such interventions are of limited success, particularly in patients where the tumor has undergone metastasis. There is a pressing need to improve the arsenal of therapies available to provide more precise and more effective treatment in a less invasive way.

10

Prostate cancer is the fastest growing neoplasm in men with an estimated 244,000 new cases in the United States being diagnosed in 1995, of which approximately 44,000 deaths will result. Prostate cancer is now the most frequently diagnosed cancer in men. Prostate cancer is latent; many men carry prostate cancer cells without overt signs of disease. It is associated with a high morbidity. Cancer metastasis to bone (late stage) is common and is almost always fatal.

15

20

25

Current treatments include radical prostatectomy, radiation therapy, hormonal ablation and chemotherapy. Unfortunately, in approximately 80% of cases, diagnosis of prostate cancer is established when the disease has already metastasized to the bones, thus limiting the effectiveness of surgical treatments. Hormonal therapy frequently fails with time with the development of hormone-resistant tumor cells. Although chemotherapeutic agents have been used in the treatment of prostate cancer, no single agent has demonstrated superiority over its counterparts, and no drug combination seems particularly effective. The generally drug-resistant, slow-growing nature of most prostate cancers makes them particularly unresponsive to standard chemotherapy.

10

15

A major, indeed the overwhelming, obstacle to cancer therapy is the problem of selectivity; that is, the ability to inhibit the multiplication of tumor cells, while leaving unaffected the function of normal cells. For example, in prostate cancer therapy, the therapeutic ratio, or ratio of tumor cell killing to normal cell killing of traditional tumor chemotherapy, is only 1.5:1. Thus, more effective treatment methods and pharmaceutical compositions for therapy and prophylaxis of neoplasia are needed.

Solid tumors frequently contain regions that are poorly vascularized, partly because the tumor cells grow faster than the endothelial cells that make up the blood vessels. Tumor cells can remain viable in such hypoxic conditions and are often refractory to chemotherapy and radiation therapy. In a recent study of cervical cancer, the oxygen status of a tumor was shown to be the single most important prognostic factor, ahead of age of patient, menopausal status, clinical stage, size and histology. Hoeckel et al. (1996) *Semin. Radiat. Oncol.* 6:1-8.

Of particular interest is development of more specific, targeted forms of cancer therapy, especially for cancers that are difficult to treat successfully. In contrast to conventional cancer therapies, which result in relatively non-specific and often serious toxicity, more specific treatment modalities attempt to inhibit or kill malignant cells selectively while leaving healthy cells intact. Radioresistant and chemoresistant tumors present particular challenges, and there is a need for methods of treating these types of tumors.

One possible treatment approach for many of these cancers is gene therapy, whereby a gene of interest is introduced into the malignant cell. Various viral vectors, including adenoviral vectors, have been developed as vehicles for gene therapy. The virtually exclusive focus in development of adenoviral vectors for gene therapy is use of adenovirus merely as a vehicle for introducing the gene of interest, not as an effector in itself. Replication of adenovirus has been viewed as an undesirable result, largely due to the host immune response. In the treatment of

25

cancer by replication-defective adenoviruses, the host immune response limits the duration of repeat doses at two levels. First, the capsid proteins of the adenovirus delivery vehicle itself are immunogenic. Second, viral late genes are frequently expressed in transduced cells, eliciting cellular immunity. Thus, the ability to repeatedly administer cytokines, tumor suppressor genes, ribozymes, suicide genes, or genes which convert prodrug to an active drug has been limited by the immunogenicity of both the gene transfer vehicle and the viral gene products of the transfer vehicle as well as the transient nature of gene expression.

10

5

Use of adenoviral vectors as therapeutic vehicles for cancer has been reported. Some of these approaches utilize tissue (i.e., cell type) specific transcriptional regulatory elements to selectively drive adenoviral replication (and thus cytotoxcity). U.S. Pat. No. 5,698,443; see also WO 95/11984; WO 96/17053; WO 96/34969; WO 98/35028. While useful and promising, there remain other treatment contexts for which tissue specific replication may be insufficient.

15

Besides cancerous cells, it is often desirable to selectively destroy certain unwanted cells or tissues. Besides surgery, however, which is invasive, there is a dearth of methods available, particularly non-invasive methods, which would allow such selective cytotoxicity and/or suppression.

20

There is a need for vector constructs that are capable of eliminating essentially all cancerous cells in a minimum number of administrations before specific immunological response against the vector prevents further treatment and which are suitable for use in specific, focused cancer ablation treatments. There is also a need for an ability to selectively destroy, or impair, unwanted cells, regardless of cell type and/or regardless of anatomical location.

10

15

#### SUMMARY OF THE INVENTION

Replication-competent adenoviral vectors specific for cells in a given, or particular, physiological state that permits or induces expression of polynucleotides under transcriptional control of a cell status-specific TRE, and methods for their use are provided. In these replication-competent adenovirus vectors, one or more adenoviral genes is under transcriptional control of an cell status-specific transcriptional regulatory element (TRE). Preferably, the adenoviral gene under transcriptional control of a cell status-specific TRE is one that is essential for adenoviral propagation. A transgene under control of the cell status-specific TRE may also be present. For the adenoviral vectors of the present invention, a cell status-specific TRE is active in a cell existing in a particular, reversible, physiological state, which may be an aberrant physiological state, i.e., a physiological state that deviates from the typical, or normal, physiological state of that same cell when in a non-dividing or regulated dividing state under normal, physiological conditions.

Accordingly, in one aspect, the invention provides an adenovirus vector comprising an adenovirus gene, wherein said adenovirus gene is under transcriptional control of a cell status-specific TRE. In another embodiment, a cell status-specific TRE is human. In another embodiment, a cell status-specific TRE comprises a cell status-specific promoter and enhancer. In yet another embodiment, a cell status-specific TRE is juxtaposed with a cell type-specific TRE, and together the two elements control expression of an adenovirus gene. Thus, the invention provides adenovirus vectors comprising a TRE comprising a cell status-specific TRE and a cell type-specific TRE.

In some embodiments, the adenovirus gene under transcriptional control of a cell status-specific TRE is an adenovirus gene essential for replication. In some embodiments, the adenoviral gene essential for replication is an early gene. In another embodiment, the early gene is E1A. In another embodiment, the early gene is E1B. In yet another embodiment, both E1A and E1B are under transcriptional

20

control of a cell status-specific TRE. In other embodiments, the adenovirus gene essential for replication is a late gene.

In another embodiment, the cell status-specific TRE comprises a hypoxia responsive element. In another embodiment, the cell status-specific TRE comprises the nucleotide sequence of SEQ ID NO:1.

In another embodiment, the cell status-specific TRE comprises a cell cycle-specific TRE. The cell cycle-specific TRE can be derived from the E2F1 5' flanking region. In one embodiment, the cell cycle-specific TRE comprises the nucleotide sequence depicted in SEQ ID NO:2.

In other embodiments, the adenovirus vector can further comprise a transgene, wherein said transgene is under transcriptional control of an cell status-specific TRE. In some embodiments, the transgene is a cytotoxic gene.

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a second adenoviral gene essential for adenoviral replication under control of a second cell status-specific TRE. The first and the second cell status-specific TREs can be identical, substantially identical, or different from, one another.

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a transgene under control of a second cell status-specific TRE. The first and the second cell status-specific TREs can be substantially identical to, or different from, one another.

In other embodiments, the adenovirus vector comprises an adenovirus gene under transcriptional control of a cell status-specific TRE, and a second adenovirus gene under transcriptional control of a cell type-specific TRE. In other embodiments, the adenovirus vector comprises an adenovirus gene under transcriptional control of a cell status-specific TRE, and a transgene under transcriptional control of a cell type-specific TRE.

15

5

10

20

In another aspect, the invention provides a host cell comprising the adenovirus vector(s) described herein.

In another aspect, the invention provides pharmaceutical compositions comprising an adenovirus vector(s) described herein.

In another aspect, the invention provides kits which contain an adenoviral vector(s) described herein.

In another aspect, methods are provided for conferring selective cytoxicity in target cells (i.e., cells which permit or induce a cell status-specific TRE to function), comprising contacting the cells with an adenovirus vector(s) described herein, whereby the vector enters the cell.

Another embodiment of the invention is an adenovirus which replicates preferentially in mammalian cells whose cell status permits or induces the function of a cell status-specific TRE.

In another aspect, methods are provided for propagating an adenovirus specific for mammalian cells whose cell status permits the function of a cell status-specific TRE, said method comprising combining an adenovirus vector(s) described herein with mammalian cells whose cell status permits the function of a cell status-specific TRE, whereby said adenovirus is propagated.

The invention further provides methods of suppressing tumor cell growth, more particularly a target tumor cell (i.e., a tumor cell that permits or induces a cell status-specific TRE to function), comprising contacting a tumor cell with an adenoviral vector of the invention such that the adenoviral vector enters the tumor cell and exhibits selective cytotoxicity for the tumor cell.

In another aspect, methods are provided for detecting cells whose cell status permits the function of a cell status-specific TRE in a biological sample, comprising contacting cells of a biological sample with an adenovirus vector(s) described herein, and detecting replication of the adenovirus vector, if any.

15

10

5

20

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of adenovirus vector CN796, in which the E1A gene is under transcriptional control of an HRE and a PSA-TRE, as described in Example 1.

5

Figure 2 shows the nucleotide sequence of an HRE from the 5' flanking region of a rat enolase-1 gene (SEQ ID NO:1).

Figure 3 shows the nucleotide sequence of the 5' flanking region of a human E2F1 gene (SEQ ID NO:2). The asterisk indicates the transcription start site.

10

Figure 4 depicts a nucleotide sequence of a prostate-specific antigen TRE.

Figure 5 depicts a nucleotide sequence of a carcinoembryonic antigen TRE.

Figure 6 depicts a nucleotide sequence of a human glandular kallikrein TRE.

Figure 7 depicts a nucleotide sequence of a mucin TRE.

Figure 8 depicts a nucleotide sequence of a rat probasin TRE.

Figure 9 depicts a nucleotide sequence and translated amino acid sequence of an adenovirus death protein.

15

#### MODES FOR CARRYING OUT THE INVENTION

20

We have discovered and constructed replication-competent adenovirus vectors which contain an adenoviral gene under transcriptional control of a cell status-specific transcriptional response element (TRE) such that the adenovirus gene is transcribed preferentially in cells whose cell status permit the function of the cell status-specific TRE, and have developed methods using these adenovirus vectors. In some preferred embodiments, the adenovirus vectors of this invention comprise at least one adenovirus gene necessary for adenoviral replication, preferably at least one early gene, under the transcriptional control of a TRE specifically regulated by binding of transcriptional factor(s) and/or co-factor(s) necessary for transcription regulated by the cell status-specific TRE. By providing for cell status-specific

25

transcription of at least one adenovirus gene required for replication, the invention

provides adenovirus vectors that can be used for specific cytotoxic effects due to selective replication and/or selective transcription. This is especially useful in the cancer context, in which targeted cell killing is desirable. This is also useful for targeted cytotoxic effects in other, non-tumor cells, when selective destruction and/or suppression of these cells is desirable. The vectors can also be useful for detecting the presence of cells whose cell status permits function of a cell status-specific TRE in, for example, an appropriate biological (such as clinical) sample. Further, the adenovirus vector(s) can optionally selectively produce one or more proteins of interest in a target cell by using a cell status-specific TRE.

10

5

We have found that adenovirus vectors of the invention replicate and/or express an adenoviral gene operably linked to a cell status-specific TRE preferentially in cells whose status permits the function of a cell status-specific TRE. In contrast to previously- described adenoviral vectors designed to replicate preferentially in specific, differentiated cell types, the adenovirus vectors of the present invention comprise regulatory elements that are not cell type-specific. Rather, they confer cell status-specific adenoviral replication and/or cell status-specific expression of an operably linked adenoviral gene and/or transgene.

The adenovirus vectors of the present invention comprise a cell status-specific

15

TRE which is functional in a cell which exhibits a particular physiological (i.e., environmental or metabolic) characteristic which is reversible and/or progressive. The target cell may exhibit an aberrant physiological state, such as low oxygen

20

tension, or may be subjected to an aberrant environmental condition, such as heat or ionizing radiation, in order for the cell-status TRE to function. The replication preference of these vectors is indicated by comparing the level of replication (i.e., titer) in cells in a requisite physiological state or condition (for example, an aberrant physiological state) to the level of replication in cells not exhibiting the requisite

25

physiological state (for example, under normal physiological conditions). Thus, the invention also uses and takes advantage of what has been considered an undesirable

10

15

20

25

aspect of adenoviral vectors, namely, their replication and possibly concomitant immunogenicity. The probability of runaway infection is significantly reduced due to the cell status-specific requirements for viral replication. Without wishing to be bound by any particular theory, the inventors note that production of adenovirus proteins can serve to activate and/or stimulate the immune system, generally and/or specifically toward target cells producing adenoviral proteins, which can be an important consideration in the cancer context, where patients are often moderately to severely immunocompromised.

The adenovirus vectors of the present invention find particular utility in specific treatment regimens, in which the treatment is highly focused toward, for example, a particular cancer which might otherwise be inoperable or untreatable. An important feature of the invention is that the vectors are useful in these treatments regardless of the tissue or cell type of the cancer, and yet their cytotoxicity can be targeted to certain locations.

#### General Techniques

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", second edition (Sanbrook et al., 1989); "Oligonucleotide Synthesis" (M.J. Gait, ed., 1984); "Animal Cell Culture" (R.I. Freshney, ed., 1987); "Methods in Enzymology" (Academic Press, Inc.); "Handbook of Experimental Immunology" (D.M. Wei & C.C. Blackwell, eds.); "Gene Transfer Vectors for Mammalian Cells" (J.M. Miller & M.P. Calos, eds., 1987); "Current Protocols in Molecular Biology" (F.M. Ausubel et al., eds., 1987); "PCR: The Polymerase Chain Reaction", (Mullis et al., eds., 1994); "Current Protocols in Immunology" (J.E. Coligan et al., eds., 1991).

For techniques related to adenovirus, see, *inter alia*, Felgner and Ringold (1989) *Nature* 337:387-388; Berkner and Sharp (1983) *Nucl. Acids Res.* 11:6003-6020; Graham (1984) *EMBO J.* 3:2917-2922; Bett et al. (1993) *J. Virology* 67:5911-5921; Bett et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:8802-8806.

#### Definitions

5

10

15

As used herein, a "transcription response element" or "transcriptional regulatory element", or "TRE" is a polynucleotide sequence, preferably a DNA sequence, which increases transcription of an operably linked polynucleotide sequence in a host cell that allows that TRE to function. A TRE can comprise an enhancer and/or a promoter.

As used herein, the term "cell status-specific TRE" is one that confers transcriptional activation on an operably linked polynucleotide in a cell which allows a cell status-specific TRE to function, i.e., a cell which exhibits a particular physiological condition, including, but not limited to, an aberrant physiological state. "Cell status" thus refers to a given, or particular, physiological state (or condition) of a cell, which is reversible and/or progressive. The physiological state may be generated internally or externally; for example, it may be a metabolic state (such as low oxygen), or it may be generated due to heat or ionizing radiation. "Cell status" is distinct from a "cell type", which relates to a differentiation state of a cell, which under normal conditions is irreversible. Generally (but not necessarily), as discussed herein, a cell status is embodied in an aberrant physiological state, examples of which are given below.

A "normal cell status" or "normal physiological state" is the status of a cell which exists in normal physiological conditions and which is non-dividing or divides in a regulated manner, i.e., a cell in a normal physiological state.

The terms "aberrant cell status" and "aberrant physiological state", used interchangeably herein, intend a condition of a cell which is a response to, a result of, or is influenced by, an aberrant physiological condition. An aberrant cell status is

25

10

15

20

25

neither cell type-specific nor tissue type-specific. An aberrant cell status is defined in relation to a cell of the same type which is in a non-dividing/regulated dividing state and under normal physiological conditions.

"Normal physiological conditions" are known to those skilled in the art.

These conditions may vary, depending on a cell's location in the body. For example, oxygen tension can vary from tissue to tissue. For *in vitro* analyses for the purposes of determining whether a TRE is responsive to deviations from normal physiological conditions, these conditions generally include an oxygen concentration of about 20% O<sub>2</sub>, and a temperature of about 37°C. "Regulated cell division" is a term well understood in the art and refers to the normal mitotic activity of a cell. Those skilled in the art understand that normal mitotic activity varies from cell type to cell type. For example, many terminally differentiated cells in tissues exhibit little or no mitotic activity, while hematopoietic cells are generally mitotically active.

An "aberrant physiological condition" or "aberrant physiological state", as used herein, intends a condition which deviates from normal physiological conditions, and includes, but is not limited to, a physiological condition that is characterized by alterations in oxygen concentration, such as hypoxic conditions; temperatures which deviate from physiological temperatures; a condition that triggers apoptosis; radiation, including ionizing radiation and UV irradiation; de-regulated cell division, resulting for example, from a lack of, or insufficient amounts of, or inactivity of, a factor which controls cell division, such as, for example, retinoblastoma protein (Rb); variations in timing of cell cycle; infection with a pathogen; exposure to a chemical substance; or a combination of the above-listed conditions. Another example is a mutation that could, or does, exist in any cell type, i.e., its existence does not depend on, or is not related to, the differentiation state of the cell.

A "target cell", as used herein, is one that permits or induces the function of a cell status-specific TRE such that it effects transcriptional activation of an operably

5

20

25

linked polynucleotide. A target cell is one which exhibits a requisite physiological (or environmental) state, which may be an aberrant physiological state. Preferably, a target cell is a mammalian cell, preferably a human cell. A target cell may or may not be neoplastic. By transcriptional activation, it is intended that transcription is increased in the target cell above the levels in a control cell (e.g., a that cell when not exhibiting a requisite physiological state (generally a normal physiological state) by at least about 2 fold, preferably at least about 5 fold, preferably at least about 10 fold, more preferably at least about 50 fold, more preferably at least about 100 fold, more preferably at least about 200 fold, even more preferably at least about 400 fold to about 500 fold, even more preferably at least about 400 fold to about 500 fold, even more preferably at least about 1000 fold. The normal levels are generally the level of activity (if any) in a cell as tested under conditions that activate the cell status-specific TRE, or the level of activity (if any) of a reporter construct lacking a cell status-specific TRE as measured in a cell exhibiting the requisite physiological condition.

A "functionally-preserved" variant of a cell status-specific TRE is a cell status-specific TRE which differs from another cell status-specific TRE, but still retains cell status cell-specific transcription activity. The difference in an cell status-specific TRE can be due to differences in linear sequence, arising from, for example, single base mutation(s), addition(s), deletion(s), and/or modification(s) of the bases. The difference can also arise from changes in the sugar(s), and/or linkage(s) between the bases of a cell status-specific TRE.

An "adenovirus vector" or "adenoviral vector" (used interchangeably) comprises a polynucleotide construct of the invention. A polynucleotide construct of this invention may be in any of several forms, including, but not limited to, DNA, DNA encapsulated in an adenovirus coat, DNA packaged in another viral or viral-like form (such as herpes simplex, and AAV), DNA encapsulated in liposomes, DNA complexed with polylysine, complexed with synthetic polycationic molecules, conjugated with transferrin, and complexed with compounds such as PEG to

immunologically "mask" the molecule and/or increase half-life, and conjugated to a nonviral protein. Preferably, the polynucleotide is DNA. As used herein, "DNA" includes not only bases A, T, C, and G, but also includes any of their analogs or modified forms of these bases, such as methylated nucleotides, internucleotide modifications such as uncharged linkages and thioates, use of sugar analogs, and modified and/or alternative backbone structures, such as polyamides. For purposes of this invention, adenovirus vectors are replication-competent in a target cell.

10

5

15

20

25

The terms "polynucleotide" and "nucleic acid", used interchangeably herein, refer to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. These terms include a single-, double- or triple-stranded DNA, genomic DNA, cDNA, RNA, DNA-RNA hybrid, or a polymer comprising purine and pyrimidine bases, or other natural, chemically, biochemically modified, non-natural or derivatized nucleotide bases. The backbone of the polynucleotide can comprise sugars and phosphate groups (as may typically be found in RNA or DNA), or modified or substituted sugar or phosphate groups. Alternatively, the backbone of the polynucleotide can comprise a polymer of synthetic subunits such as phosphoramidates and thus can be a oligodeoxynucleoside phosphoramidate (P-NH2) or a mixed phosphoramidate-phosphodiester oligomer. Peyrottes et al. (1996) Nucleic Acids Res. 24: 1841-8; Chaturvedi et al. (1996) Nucleic Acids Res. 24: 2318-23; Schultz et al. (1996) Nucleic Acids Res. 24: 2966-73. A phosphorothiate linkage can be used in place of a phosphodiester linkage. Braun et al. (1988) J. Immunol. 141: 2084-9; Latimer et al. (1995) Mol. Immunol. 32: 1057-1064. In addition, a double-stranded polynucleotide can be obtained from the single stranded polynucleotide product of chemical synthesis either by synthesizing the complementary strand and annealing the strands under appropriate conditions, or by synthesizing the complementary strand de novo using a DNA polymerase with an appropriate primer.

fragment, exons, introns, mRNA, tRNA, rRNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs, uracyl, other sugars and linking groups such as fluororibose and thioate, and nucleotide branches. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. Other types of modifications included in this definition are caps, substitution of one or more of the naturally occurring nucleotides with an analog, and introduction of means for attaching the polynucleotide to proteins, metal ions, labeling components, other polynucleotides, or a solid support. Preferably, the polynucleotide is DNA. As used herein, "DNA" includes not only bases A, T, C, and G, but also includes any of their analogs or modified forms of these bases, such as methylated nucleotides, internucleotide modifications such as uncharged linkages and thioates, use of sugar analogs, and modified and/or alternative backbone structures, such as polyamides.

The following are non-limiting examples of polynucleotides: a gene or gene

20

5

10

15

25

A polynucleotide or polynucleotide region has a certain percentage (for example, 80%, 85%, 90%, or 95%) of "sequence identity" to another sequence means that, when aligned, that percentage of bases are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in *Current Protocols in Molecular Biology* (F.M. Ausubel et al., eds., 1987) Supplement 30, section 7.7.18, Table 7.7.1. A preferred alignment program is ALIGN Plus (Scientific and Educational Software, Pennsylvania), preferably using default parameters.

"Under transcriptional control" is a term well-understood in the art and indicates that transcription of a polynucleotide sequence, usually a DNA sequence,

10

15

20

depends on its being operably (operatively) linked to an element which contributes to the initiation of, or promotes, transcription. "Operably linked" refers to a juxtaposition wherein the elements are in an arrangement allowing them to function.

"Replication" and "propagation" are used interchangeably and refer to the ability of a polynucleotide construct of the invention to reproduce, or proliferate. This term is well understood in the art. For purposes of this invention, replication involves production of adenovirus proteins and is generally directed to reproduction of adenovirus. Replication can be measured using assays standard in the art and described herein, such as a burst assay, plaque assay, or a one-step growth curve assay.

As used herein, "cytotoxicity" is a term well understood in the art and refers to a state in which a cell's usual biochemical or biological activities are compromised (i.e., inhibited). These activities include, but are not limited to, metabolism; cellular replication; DNA replication; transcription; translation; uptake of molecules. "Cytotoxicity" includes cell death and/or cytolysis. Assays are known in the art which indicate cytotoxicity, such as dye exclusion, <sup>3</sup>H-thymidine uptake, and plaque assays.

The term "selective cytotoxicity", as used herein, refers to the cytotoxicity conferred by an adenovirus vector of the present invention on a cell which allows or induces a cell status-specific TRE to function (a target cell) when compared to the cytotoxicity conferred by an adenoviral vector of the present invention on a cell which does not allow a cell status-specific TRE to function (a non-target cell). Such cytotoxicity may be measured, for example, by plaque assays, by reduction or stabilization in size of a tumor comprising target cells, or the reduction or stabilization of serum levels of a marker characteristic of the tumor cells, or a tissue-specific marker, e.g., a cancer marker, such as prostate specific antigen.

In the context of adenovirus, a "heterologous polynucleotide" or "heterologous gene" or "transgene" is any polynucleotide or gene that is not present

10

15

20

25

in wild-type adenovirus. Preferably, the transgene will also not be expressed or present in the target cell prior to introduction by the adenovirus vector. Examples of preferred transgenes are provided below.

In the context of adenovirus, a "heterologous" promoter or enhancer is one which is not associated with or derived from an adenovirus gene.

In the context of adenovirus, an "endogenous" promoter, enhancer, or TRE is native to or derived from adenovirus.

In the context of a cell status-specific TRE, a "heterologous" promoter or enhancer is one which is not normally associated in a cell with or derived from a cell status-specific TRE. Examples of a heterologous promoter or enhancer are the albumin promoter or enhancer and other viral promoters and enhancers, such as SV40, or cell type-specific TREs such as a prostate-specific TRE.

A "cell type-specific TRE" is preferentially functional in a specific type of cell relative to other types of cells. In contrast to cell status, "cell type" is a reflection of a differentiation state of a cell which is irreversible. For example, a prostate-specific antigen is expressed in prostate cells, but is not substantially expressed in other cell types such as hepatocytes, astrocytes, cardiocytes, lymphocytes, etc. Generally, a cell type-specific TRE is active in only one cell type. When a cell type-specific TRE is active in more than one cell type, its activity is restricted to a limited number of cell types, i.e., it is not active in all cell types. A cell type-specific TRE may or may not be tumor cell specific.

"Suppressing" tumor growth indicates a growth state that is curtailed when compared to growth without contact with, i.e., transfection by, an adenoviral vector described herein. Tumor cell growth can be assessed by any means known in the art, including, but not limited to, measuring tumor size, determining whether tumor cells are proliferating using a <sup>3</sup>H-thymidine incorporation assay, or counting tumor cells. "Suppressing" tumor cell growth means any or all of the following states: slowing, delaying, and stopping tumor growth, as well as tumor shrinkage.

10

15

20

As used herein, the terms "neoplastic cells", "neoplasia", "tumor", "tumor cells", "cancer" and "cancer cells", (used interchangeably) refer to cells which exhibit relatively autonomous growth, so that they exhibit an aberrant growth phenotype characterized by a significant loss of control of cell proliferation (i.e., de-regulated cell division). Neoplastic cells can be malignant or benign.

A "host cell" includes an individual cell or cell culture which can be or has been a recipient of an adenoviral vector(s) of this invention. Host cells include progeny of a single host cell, and the progeny may not necessarily be completely identical (in morphology or in total DNA complement) to the original parent cell due to natural, accidental, or deliberate mutation and/or change. A host cell includes cells transfected or infected *in vivo* or *in vitro* with an adenoviral vector of this invention.

"Replication" and "propagation" are used interchangeably and refer to the ability of an adenovirus vector of the invention to reproduce or proliferate. These terms are well understood in the art. For purposes of this invention, replication involves production of adenovirus proteins and is generally directed to reproduction of adenovirus. Replication can be measured using assays standard in the art and described herein, such as a burst assay or plaque assay. "Replication" and "propagation" include any activity directly or indirectly involved in the process of virus manufacture, including, but not limited to, viral gene expression; production of viral proteins, nucleic acids or other components; packaging of viral components into complete viruses; and cell lysis.

An "ADP coding sequence" is a polynucleotide that encodes ADP or a functional fragment thereof. In the context of ADP, a "functional fragment" of ADP is one that exhibits cytotoxic activity, especially cell lysis, with respect to adenoviral replication. Ways to measure cytotoxic activity are known in the art and are described herein.

A polynucleotide that "encodes" an ADP polypeptide is one that can be transcribed and/or translated to produce an ADP polypeptide or a fragment thereof. The anti-sense strand of such a polynucleotide is also said to encode the sequence.

An "ADP polypeptide" is a polypeptide containing at least a portion, or region, of the amino acid sequence of an ADP (see, for example, SEQ ID NO:5), and which displays a function associated with ADP, particularly cytotoxicity, more particularly, cell lysis. As discussed herein, these functions can be measured using techniques known in the art. It is understood that certain sequence variations may be used, due to, for example, conservative amino acid substitutions, which may provide ADP polypeptides.

A polynucleotide sequence that is "depicted in" a SEQ ID NO means that the sequence is present as an identical contiguous sequence in the SEQ ID NO. The term encompasses portions, or regions of the SEQ ID NO as well as the entire sequence contained within the SEQ ID NO.

A "biological sample" encompasses a variety of sample types obtained from an individual and can be used in a diagnostic or monitoring assay. The definition encompasses blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom, and the progeny thereof. The definition also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as proteins or polynucleotides. The term "biological sample" encompasses a clinical sample, and also includes cells in culture, cell supernatants, cell lysates, serum, plasma, biological fluid, and tissue samples.

An "individual" is a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, farm animals, sport animals, rodents, primates, and pets.

An "effective amount" is an amount sufficient to effect beneficial or desired results, which may include clinical results. An effective amount can be administered

15

5

10

20

in one or more administrations. For purposes of this invention, an effective amount of an adenoviral vector is an amount that is sufficient to palliate, ameliorate, stabilize, reverse, slow or delay the progression of the disease state.

#### Adenoviral vectors comprising a cell status-specific TRE

10

5

15

20

25

The present invention provides adenoviral vector constructs which comprise an adenovirus gene under transcriptional control of a cell status-specific TRE. Preferably, the adenovirus gene contributes to cytotoxicity (whether direct and/or indirect), more preferably one that contributes to or causes cell death, even more preferably is essential for advenoviral replication. Examples of a gene that contributes to cytotoxicity include, but are not limited to, adenovirus death protein (ADP; discussed below). When the adenovirus vector(s) is selectively (i.e., preferentially) replication competent for propagation in target cells, i.e., cells which permit or induce a cell-status TRE to function, these cells will be preferentially killed upon adenoviral proliferation. Once the target cells are destroyed due to selective cytotoxic and/or cytolytic replication, the adenovirus vector replication is significantly reduced, thus lessening the probability of runaway infection and undesirable bystander effects. In vitro cultures may be retained to monitor the mixture (such as, for example, a biopsy or other appropriate biological sample) for occurrence (i.e., presence) and/or recurrence of the target cell, e.g., a neoplastic cell or other undesired cell. To further ensure cytotoxicity, one or more transgenes having a cytotoxic effect may also be present and under selective transcriptional control. In this embodiment, one may provide higher confidence that the target cells will be destroyed. Additionally, or alternatively, an adenovirus gene that contributes to cytotoxicity and/or cell death (such as ADP) may be included in the adenoviral vector, either free of, or under, selective transcriptional control.

Cell status-specific TREs for use in the adenoviral vectors of the present invention can be derived from any species, preferably a mammal. A number of genes have been described which are expressed in response to, or in association with, a cell

10

15

20

status. Any of these cell status-associated genes may be used to generate a cell status-specific TRE.

An example of a cell status is cell cycle. An exemplary gene whose expression is associated with cell cycle is E2F-1, a ubiquitously expressed, growth-regulated gene, which exhibits peak transcriptional activity in S phase. Johnson et al. (1994) *Genes Dev.* 8:1514-1525. The RB protein, as well as other members of the RB family, form specific complexes with E2F-1, thereby inhibiting its ability to activate transcription. Thus, E2F-1-responsive promoters are down-regulated by RB. Many tumor cells have disrupted RB function, which can lead to de-repression of E2F-1-responsive promoters, and, in turn, de-regulated cell division.

Accordingly, in one embodiment, the invention provides an adenoviral vector in which an adenoviral gene (preferably a gene necessary for replication) is under transcriptional control of a cell status-specific TRE, wherein the cell status-specific TRE comprises a cell cycle-activated, or cell-cycle specific, TRE. In one embodiment, the cell cycle-activated TRE is an E2F1 TRE. In one embodiment, this TRE comprises the sequence depicted in Figure 3 and SEQ ID NO:2.

Another group of genes which are regulated by cell status are those whose expression is increased in response to hypoxic conditions. Bunn and Poyton (1996) *Physiol. Rev.* 76:839-885; Dachs and Stratford (1996) *Br. J. Cancer* 74:5126-5132; Guillemin and Krasnow (1997) *Cell* 89:9-12. Many tumors have insufficient blood supply, due in part to the fact that tumor cells typically grow faster than the endothelial cells that make up the blood vessels, resulting in areas of hypoxia in the tumor. Folkman (1989) *J. Natl. Cancer Inst.* 82:4-6; and Kallinowski (1996) *The Cancer J.* 9:37-40. An important mediator of hypoxic responses is the transcriptional complex HIF-1, or hypoxia inducible factor-1, which interacts with a hypoxia-responsive element (HRE) in the regulatory regions of several genes, including vascular endothelial growth factor, and several genes encoding glycolytic enzymes, including enolase-1. Murine HRE sequences have been identified and characterized.

Firth et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:6496-6500. An HRE from a rat enolase-1 promoter is described in Jiang et al. (1997) *Cancer Res.* 57:5328-5335. An HRE from a rat enolase-1 promoter is depicted in Figure 2 and given as SEQ ID NO:1.

5

Accordingly, in one embodiment, an adenovirus vector comprises an adenovirus gene, preferably an adenoviral gene essential for replication, under transcriptional control of a cell status-specific TRE comprising an HRE. In one embodiment, the cell status-specific TRE comprises the HRE depicted in Figure 2 and SEQ ID NO:1.

10

15

Other cell status-specific TREs include heat-inducible (i.e., heat shock) promoters, and promoters responsive to radiation exposure, including ionizing radiation and UV radiation. For example, the promoter region of the early growth response-1 (Egr-1) gene contains an element(s) inducible by ionizing radiation. Hallahan et al. (1995) *Nat. Med.* 1:786-791; and Tsai-Morris et al. (1988) *Nucl. Acids. Res.* 16:8835-8846. Heat-inducible promoters, including heat-inducible elements, have been described. See, for example Welsh (1990) in "Stress Proteins in Biology and Medicine", Morimoto, Tisseres, and Georgopoulos, eds. Cold Spring Harbor Laboratory Press; and Perisic et al. (1989) *Cell* 59:797-806. Accordingly, in some embodiments, the cell status-specific TRE comprises an element(s) responsive to ionizing radiation. In one embodiment, this TRE comprises a 5' flanking sequence of an Egr-1 gene. In other embodiments, the cell status-specific TRE comprises a heat shock responsive, or heat-inducible, element.

20

A cell status-specific TRE can also comprise multimers. For example, an HRE can comprise a tandem series of at least two, at least three, at least four, or at least five hypoxia-responsive elements. These multimers may also contain heterologous promoter and/or enhancer sequences.

25

A cell status-specific TRE may or may not lack a silencer. The presence of a silencer (i.e., a negative regulatory element) may assist in shutting off transcription

10

15

20

25

(and thus replication) in non-permissive cells (i.e., cell in a normal cell state). Thus, presence of a silencer may confer enhanced cell status-specific replication by more effectively preventing adenoviral vector replication in non-target cells. Alternatively, lack of a silencer may assist in effecting replication in target cells, thus conferring enhanced cell status-specific replication due to more effective replication in target cells.

In other embodiments, the adenoviral vector comprises an adenoviral gene

essential for adenoviral replication under control of a first cell status-specific TRE, and a second adenoviral gene essential for adenoviral replication under control of a second cell status-specific TRE. The first and the second cell status-specific TREs may or may not be identical, and may or may not be substantially identical to one another. By "substantially identical" is meant a requisite degree of sequence identity between the two TREs. The degree of sequence identity between these TREs is at least about 80%, preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95%, even more preferably at least about 98%, and most preferably 100%. Sequence identity can be determined by a sequence comparison using, i.e., sequence alignment programs that are known in the art, such as those described in Current Protocols in Molecular Biology (F.M. Ausubel et al., eds., 1987) Supplement 30, section 7.7.18, Table 7.7.1 A preferred alignment program is ALIGN Plus (Scientific and Educational Software, Pennsylvania), preferably using default parameters. Alternatively, hybridization under stringent conditions can also indicate degree of sequence identity. Stringent conditions are known in the art; an example of a stringent condition is 80°C (or higher temperature) and 6 X SSC (or less concentrated SSC). Other hybridization conditions and parameters (in order of increasing stringency) are: incubation temperatures of 25°C, 37°C, 50°C, and 68°C; buffer concentrations of 10 X SSC, 6 X SSC, 1 X SSC, 0.1 X SSC (where 1 X SSC is 0.15 M NaCl and 15 mM citrate buffer) and their equivalents

using other buffer systems; formamide concentrations of 0%, 25%, 50%, and 75%;

10

15

20

incubation times from about 24 hours about 5 minutes; 1, 2, or more washing steps; wash incubation times of 1, 2, or 15 minutes; and wash solutions of 6 X SSC, 1 X SSC, 0.1 X SSC, or deionized water.

Adenoviral constructs in which the first and second cell status-specific TREs are identical or substantially identical, particularly if these TREs control transcription of early genes (such as E1A and E1B), may display an instability which may be desirable in certain contexts, such as when an automatic "self-destruction" property can shut down the virus, thereby controlling the degree of propagation. Accordingly, in some embodiments, the first and second cell status-specific TRE, or the first and second TRE (one of which is a cell-status-specific TRE) are sufficiently identical to confer instability when compared to two TREs which are less identical with respect to each other (i.e., have more sequence divergence or dissimilarity). Preferred embodiments are those in which the two TREs control E1A and E1B respectively. "Instability" means that the structural integrity of the adenoviral vectors is not preserved as the virus replicates in cells, and can be measured using standard methods in the art, such as Southern analysis. In other embodiments, the first and second TREs are sufficiently divergent and/or placed in the vector such that the vector is stable (i.e., the structural integrity of the adenoviral vector is preserved).

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a transgene under control of a second cell status-specific TRE. The first and the second cell status-specific TREs may or may not be substantially identical to one another.

In some embodiments, a cell status-specific TRE can be juxtaposed with another TRE, such as a different cell status-specific TRE, or, alternatively, a cell type-specific TRE. "Juxtaposed" means a cell status-specific TRE and the second TRE transcriptionally control the same gene, or at least are proximate with respect to the same gene. For these embodiments, the cell status-specific TRE and the second

10

15

20

25

TRE may be in any of a number of configurations, including, but not limited to, (a) next to each other (i.e., abutting); (b) both 5' to the gene that is transcriptionally controlled (i.e., may have intervening sequences between them); (c) one TRE 5' and the other TRE 3' to the gene. For example, as described in Example 1 and shown in Figure 1, a cell type-specific TRE can be juxtaposed with a cell status-specific TRE to control transcription of an operably linked adenoviral gene. Such "composite" TREs can be used to confer both cell status- and cell type-specific expression of an operably linked polynucleotide, and thus may confer significantly greater specificity and/or efficacy. Examples of cell type-specific TREs are provided below. Alternatively, "composite" TREs can be used to confer different, and possibly synergistic, cell status specificity. For example, a composite TRE could confer specificity to hypoxia and heat shock.

Example 1 provides a description of an adenovirus construct in which a composite TRE upstream of E1A consisting of an HRE and a prostate-specific TRE. PSA-TRE (which consists of enhancer sequences -5322 to -3738 fused to PSA promoter sequence -541 to +12; see U.S. Pat. Nos. 5,871,726; 5,648,478). Accordingly, in some embodiments, the invention provides an adenovirus vector comprising an adenovirus gene essential for replication, preferably an early gene, preferably E1A or E1B, under transcriptional control of a TRE comprising an HRE (preferably comprising or consisting of the 67-base fragment depicted in SEQ ID NO:1) and a prostate cell specific TRE, preferably comprising a PSA enhancer (preferably -5322 to -3738; or about 503 to about 2086 of SEQ ID NO:3 (bases about 503 to about 2086 of Figure 4), and a promoter, preferably comprising a PSA enhancer and a PSA promoter (about 5285 to about 5836 of SEQ ID NO:3).

As is readily appreciated by one skilled in the art, a cell status-specific TRE is a polynucleotide sequence, and, as such, can exhibit function over a variety of sequence permutations. Methods of nucleotide substitution, addition, and deletion are known in the art, and readily available functional assays (such as the CAT or

luciferase reporter gene assay) allow one of ordinary skill to determine whether a sequence variant exhibits requisite cell status-specific transcription function. Hence, the invention also includes functionally-preserved variants of the nucleic acid sequences disclosed herein, which include nucleic acid substitutions, additions, and/or deletions. While not wishing to be bound by a single theory, the inventors note that it is possible that certain modifications will result in modulated resultant expression levels, including enhanced expression levels. Achievement of modulated resultant expression levels, preferably enhanced expression levels, may be especially desirable in the case of certain, more aggressive forms of cancer, or when a more rapid and/or aggressive pattern of cell killing is warranted (due to an immunocompromised condition of the individual, for example).

15

10

5

20

25

As an example of how cell status-specific TRE activity can be determined, a polynucleotide sequence or set of such sequences can be generated using methods known in the art, such as chemical synthesis, site-directed mutagenesis, PCR, and/or recombinant methods. The sequence(s) to be tested is inserted into a vector containing an appropriate reporter gene, including, but not limited to. chloramphenicol acetyl transferase (CAT), β-galactosidase (encoded by the lacZ gene), luciferase (encoded by the luc gene), green fluorescent protein, alkaline phosphatase, and horse radish peroxidase. Such vectors and assays are readily available, from, inter alia, commercial sources. Plasmids thus constructed are transfected into a suitable host cell to test for expression of the reporter gene as controlled by the putative cell status-specific TRE using transfection methods known in the art, such as calcium phosphate precipitation, electroporation, liposomes (lipofection), and DEAE-dextran. Suitable host cells include any cell type, including but not limited to, Hep3B, Hep G2, HuH7, HuH1/Cl2, LNCaP, HBL-100, Chang liver cells, MCF-7, HLF, HLE, 3T3, HUVEC, and HeLa. Host cells transfected with the TRE-reporter gene construct to be tested are subjected to conditions which result in a change in cell status (for example, one which result in an aberrant physiological

10

15

20

25

state). The same cells not subjected to these conditions, i.e., which are under normal physiological conditions and therefore in a normal physiological state, serve as controls. Results are obtained by measuring the level of expression of the reporter gene using standard assays. Comparison of expression between cells in a particular state and control indicates presence or absence of transcriptional activation. "Transcriptional activation" has been defined above.

Comparisons between or among various cell status-specific TREs can be assessed, for example, by measuring and comparing levels of expression within a single cell line under normal and aberrant physiological conditions. These comparisons may also be made by measuring and comparing levels of expression within a single cell line subjected to reversible environmental conditions (such as heat) and cells not subjected to such conditions. It is understood that absolute transcriptional activity of an cell status-specific TRE will depend on several factors, such as the nature of the target cell, delivery mode and form of the cell status-specific TRE, and the coding sequence that is to be selectively transcriptionally activated. To compensate for various plasmid sizes used, activities can be expressed as relative activity per mole of transfected plasmid. Alternatively, the level of transcription (i.e., mRNA) can be measured using standard Northern analysis and hybridization techniques. Levels of transfection (i.e., transfection efficiencies) are measured by cotransfecting a plasmid encoding a different reporter gene under control of a different TRE, such as the cytomegalovirus (CMV) immediate early promoter. This analysis can also indicate negative regulatory regions, i.e., silencers.

As an example of how hypoxia induction can be measured, one can use an assay such as that described in Jiang et al. (1997) *Cancer Research* 57:5328-5335 or Dachs et al. (1997) *Nature Med.* 3:515-520. For example, a construct comprising a putative HRE, or multiple tandem copies thereof, together with a minimal promoter element, operably linked and controlling transcription of a polynucleotide which encodes a protein which is detectable or can be used to give a detectable signal, is

introduced into host cells. The host cells are then subjected to conditions of normoxia (e.g., 20% O<sub>2</sub>), and varying degrees of hypoxia, such as 5%, 2%, 1%, 0.1%, or less, O<sub>2</sub>. The expression product of the operably linked polynucleotide (reporter gene) is then measured.

5

Alternatively a putative cell status-specific TRE can be assessed for its ability to confer adenoviral replication preference for cells exhibiting the requisite physiological state, such as heat or ionizing radiation. For this assay, constructs containing an adenovirus gene essential to replication operably linked to a putative cell status-specific TRE are transfected into cells which exhibit the requisite physiological state. Viral replication in those cells is compared, for example, to viral replication by the construct in cells under normal physiological conditions (i.e., not exhibiting the requisite physiological state).

10

Any of the various serotypes of adenovirus can be used, such as Ad2, Ad5, Ad12 and Ad40. For purposes of illustration, serotype Ad5 will be exemplified herein.

15

When a cell status-specific TRE is used with an adenovirus gene that is essential for propagation replication competence is preferentially achievable in the target cell expressing cell status. Preferably, the gene is an early gene, such as E1A, E1B, E2, or E4. (E3 is not essential for viral replication.) More preferably, the early gene under cell status-TRE control is E1A and/or E1B. More than one early gene can be placed under control of an cell status-specific TRE. Example 1 provides a more detailed description of such constructs.

20

The E1A gene is expressed immediately after viral infection (0-2 hours) and before any other viral genes. E1A protein acts as a *trans*-acting positive-acting transcriptional regulatory factor, and is required for the expression of the other early viral genes E1B, E2, E3, E4, and the promoter-proximal major late genes. Despite the nomenclature, the promoter proximal genes driven by the major late promoter are expressed during early times after Ad5 infection. Flint (1982) *Biochem. Biophys*.

Acta 651:175–208; Flint (1986) Advances Virus Research 31:169–228; Grand (1987) Biochem. J. 241:25–38. In the absence of a functional E1A gene, viral infection does not proceed, because the gene products necessary for viral DNA replication are not produced. Nevins (1989) Adv. Virus Res. 31:35–81. The transcription start site of Ad5 E1A is at 498 and the ATG start site of the E1A protein is at 560 in the virus genome.

The E1B protein functions *in trans* and is necessary for transport of late mRNA from the nucleus to the cytoplasm. Defects in E1B expression result in poor expression of late viral proteins and an inability to shut off host cell protein synthesis. The promoter of E1B has been implicated as the defining element of difference in the host range of Ad40 and Ad5: clinically Ad40 is an enterovirus, whereas Ad5 causes acute conjunctivitis. Bailey, Mackay et al. (1993) *Virology* 193:631; Bailey et al. (1994) *Virology* 202:695-706). The E1B promoter of Ad5 consists of a single high-affinity recognition site for Spl and a TATA box.

The E2 region of adenovirus codes for proteins related to replication of the adenoviral genome, including the 72 kDa DNA-binding protein, the 80 kD precursor terminal protein and the viral DNA polymerase. The E2 region of Ad5 is transcribed in a rightward orientation from two promoters, termed E2 early and E2 late, mapping at 76.0 and 72.0 map units, respectively. While the E2 late promoter is transiently active during late stages of infection and is independent of the E1A transactivator

protein, the E2 early promoter is crucial during the early phases of viral replication.

The E2 late promoter overlaps with the coding sequences of a gene encoded by the counterstrand and is therefore not amenable to genetic manipulation. However, the E2 early promoter overlaps only for a few base pairs with sequences coding for a 33 kD protein on the counterstrand. Notably, the SpeI restriction site (Ad5 position 27082) is part of the stop codon for the above mentioned 33 kD protein and conveniently separates the major E2 early transcription initiation site and TATA-binding protein site from the upstream transcription factor biding sites E2F and ATF.

20

5

10

15

10

15

20

Therefore, insertion of a cell status-TRE having SpeI ends into the SpeI site in the +-strand would disrupt the endogenous E2 early promoter of Ad5 and should allow cell status-restricted expression of E2 transcripts.

The E4 gene has a number of transcription products. The E4 region codes for two polypeptides which are responsible for stimulating the replication of viral genomic DNA and for stimulating late gene expression. The protein products of open reading frames (ORFS) 3 and 6 can both perform these functions by binding the 55kD protein from E1B and heterodimers of E2F-1 and DP-1. The ORF 6 protein requires interaction with the EIB 55kD protein for activity while the ORF 3 protein does not. In the absence of functional protein from ORF 3 and ORF 6, plaques are produced with an efficiency less than 10<sup>-6</sup> that of wild type virus. To further restrict viral replication to cells exhibiting a requisite physiological condition or state, E4 ORFs 1-3 can be deleted, making viral DNA replication and late gene synthesis dependent on E4 ORF 6 protein. By combining such a mutant with sequences in which the E1B region is regulated by a cell status-specific TRE, a virus can be obtained in which both the EIB function and E4 function are dependent on a cell status-specific TRE driving EIB.

The major late genes relevant to the subject invention are genes L1, L2, L3, L4, and L5 which encode proteins of the adenovirus virion. All of these genes (typically coding for structural proteins) are probably required for adenoviral replication. The late genes are all under the control of the major late promoter (MLP), which is located in Ad5 at +5986 to +6048.

In addition to conferring selective cytotoxic and/or cytolytic activity by virtue of preferential replication competence in cells exhibiting a requisite physiological state (for example, an aberrant physiological state such as low oxygen conditions), the adenovirus vectors of this invention can further include a heterologous gene (transgene) under the control of a cell status-specific TRE. In this way, various genetic capabilities may be introduced into target cells, particularly cancer cells. For

example, in certain instances, it may be desirable to enhance the degree and/or rate of cytotoxic activity, due to, for example, the relatively refractory nature or particular aggressiveness of the cancerous target cell. This could be accomplished by coupling the cell status-specific replicative cytotoxic activity with cell-specific expression of, for example, HSV-tk and/or cytosine deaminase (cd), which renders cells capable of metabolizing 5-fluorocytosine (5-FC) to the chemotherapeutic agent 5-fluorouracil (5-FU). Using these types of transgenes may also confer a bystander effect.

Other desirable transgenes that may be introduced via an adenovirus vector(s)

10

5

include genes encoding cytotoxic proteins, such as the A chains of diphtheria toxin, ricin or abrin (Palmiter et al. (1987) *Cell* 50: 435; Maxwell et al. (1987) *Mol. Cell. Biol.* 7: 1576; Behringer et al. (1988) *Genes Dev.* 2: 453; Messing et al. (1992) *Neuron* 8: 507; Piatak et al. (1988) *J. Biol. Chem.* 263: 4937; Lamb et al. (1985) *Eur. J. Biochem.* 148: 265; Frankel et al. (1989) *Mol. Cell. Biol.* 9: 415), genes encoding a factor capable of initiating apoptosis, sequences encoding antisense transcripts or ribozymes, which among other capabilities may be directed to mRNAs encoding proteins essential for proliferation, such as structural proteins, or transcription factors; viral or other pathogenic proteins, where the pathogen proliferates intracellularly; genes that encode an engineered cytoplasmic variant of a nuclease (e.g. RNase A) or

20

15

gene, and the like. Other genes of interest include cytokines, antigens, transmembrane proteins, and the like, such as IL-1, -2, -6, -12, GM-CSF, G-CSF, M-CSF, IFN- $\alpha$ , - $\beta$ , - $\gamma$ , TNF- $\alpha$ , - $\beta$ , TGF- $\alpha$ , - $\beta$ , NGF, and the like. The positive effector genes could be used in an earlier phase, followed by cytotoxic activity due to replication.

protease (e.g. awsin, papain, proteinase K, carboxypeptidase, etc.), or encode the Fas

25

In one embodiment, the adenovirus death protein (ADP), encoded within the E3 region, is maintained in the adenovirus vector. The ADP gene, under control of the major late promoter (MLP), appears to code for a protein (ADP) that is important in expediting host cell lysis. Tollefson et al. (1996) *J. Virol.* 70(4):2296; Tollefson et

al. (1992) *J. Virol.* 66(6):3633. Thus, adenoviral vectors containing the ADP gene may render the adenoviral vector more potent, making possible more effective treatment and/or a lower dosage requirement.

that further includes a polynucleotide sequence encoding an ADP. A DNA sequence

encoding an ADP and the amino acid sequence of an ADP are depicted Figure 9.

Briefly, an ADP coding sequence is obtained preferably from Ad2 (since this is the

strain in which ADP has been more fully characterized) using techniques known in

the art, such as PCR. Preferably, the Y leader (which is an important sequence for

correct expression of late genes) is also obtained and ligated to the ADP coding

sequence. The ADP coding sequence (with or without the Y leader) can then be

be inserted in other locations of the adenovirus genome, such as the E4 region.

introduced into the adenoviral genome, for example, in the E3 region (where the ADP

coding sequence will be driven by the MLP). The ADP coding sequence could also

Alternatively, the ADP coding sequence could be operably linked to a heterologous

promoter (with or without enhancer(s)), including, but not limited to, another viral

promoter, a cell status-specific TRE such as a hypoxia responsive element, or a cell

type-specific TRE such as those derived from carcinoembryonic antigen (CEA),

Accordingly, the invention provides an adenoviral vector as described herein

5

10

15

15

20

Adenoviral vectors of the invention further comprising a cell type specific element

25

In addition to conferring selective cytotoxic and/or cytolytic activity by virtue of preferential replication competence and/or by preferential transcription of a gene encoding a cytotoxic factor in cells exhibiting a requisite physiological state, the adenovirus vectors of this invention can further include an adenovirus gene and/or a heterologous gene (transgene) under the control of a cell type-specific TRE. In this way, cytotoxicity is further limited to a particular cell type.

mucin, and rat probasin genes.

For example, TREs that function preferentially in prostate cells include, but are not limited to, TREs derived from the prostate-specific antigen gene (*PSA*-TRE) (U.S. Patent No. 5,648,478), the glandular kallikrein-1 gene (from the human gene, *hKLK2*-TRE), and the probasin gene (*PB*-TRE) (International Patent Application No. PCT/US98/04132). All three of these genes are preferentially expressed in prostate cells and the expression is androgen-inducible. Generally, expression of genes responsive to androgen induction requires the presence of an androgen receptor (AR).

PSA is synthesized exclusively by normal, hyperplastic, and malignant prostatic epithelia; hence, its tissue-specific expression has made it an excellent biomarker for benign prostatic hyperplasia (BPH) and prostatic carcinoma (CaP). Normal serum levels of PSA are typically below 5 ng/ml, with elevated levels indicative of BPH or CaP. Lundwall et al. (1987) *FEBS Lett.* 214: 317; Lundwall (1989) *Biochem. Biophys. Res. Comm.* 161: 1151; and Riegmann et al. (1991) *Molec. Endocrin.* 5: 1921.

15

20

10

5

The region of the *PSA* gene that is used to provide cell specificity dependent upon androgens, particular in prostate cells, involves approximately 6.0 kilobases. Schuur et al. (1996) *J. Biol. Chem.* 271:7043-7051. An enhancer region of approximately 1.5 kb in humans is located between nt -5322 and nt -3738, relative to the transcription start site of the *PSA* gene. The *PSA* promoter consists of the sequence from about nt -540 to nt +12 relative to the transcription start site. Juxtapositioning of these two genetic elements yield a fully functional, minimal prostate-specific enhancer/promoter (*PSE*) TRE. Other portions of the approximately 6.0 kb region of the *PSA* gene can be used in the present invention, as long as requisite functionality is maintained. In Example 1, adenoviral vector CN796 is described which comprises a composite TRE comprising an HRE and a PSA-TRE, the PSA-TRE comprising a PSA enhancer from -5322 to -3738 fused to a PSA promoter from -541 to +12. This PSA-TRE is derived from adenoviral vector CN706. Rodriguez et al. (1997) *Cancer Research* 57:2559-2563. Accordingly, in

one embodiment an adenoviral vector comprises and adenovirus E1A gene under transcriptional control of a composite TRE comprising the cell status-specific TRE, HRE, and a cell type-specific TRE, a PSA-TRE.

5

10

15

20

25

The *PSE* and *PSA* TRE used in the present invention are derived from sequences depicted in Figure 4 (SEQ ID NO:3). The enhancer element is nucleotides about 503 toabout 2086 of Figure 4 (SEQ ID NO:3). The promoter is nucleotides about 5285 to about 5836 of Figure 4 (SEQ ID NO:3). Accordingly, in some embodiments, the composite TRE comprises an HRE comprising SEQ ID NO:1 and a PSA-TRE comprises nucleotides about 503 to about 2086 of SEQ ID NO:3. In other embodiments, the composite TRE comprises an HRE comprising SEQ ID NO:1 and a PSA-TRE comprises nucleotides about 503 to about 2086 of SEQ ID NO:3 and nucleotides about 5285 to about 5836 of SEQ ID NO:3. As described above, these composite (HRE/PSA) TREs may be operably linked to an adenovirus gene essential for replication, especially an early gene such as E1A or E1B. Example 1 describes such a construct.

In the present invention, replication-competent adenovirus vectors comprising a cell status-specific TRE and a cell type-specific TRE may employ cell type-specific TREs that are preferentially functional in particular tumor cells. Non-limiting examples of tumor cell-specific TREs, and non-limiting examples of respective potential target cells, include TREs from the following genes: α-fetoprotein (*AFP*) (liver cancer), mucin-like glycoprotein DF3 (*MUC1*) (breast carcinoma), carcinoembryonic antigen (*CEA*) (colorectal, gastric, pancreatic, breast, and lung cancers), plasminogen activator urokinase (*uPA*) and its receptor gene (breast, colon, and liver cancers), *HER-2/neu* (*c-erbB2/neu*) (breast, ovarian, stomach, and lung cancers).

Other cell type-specific TREs may be derived from the following exemplary genes (cell type in which the TREs are specifically functional are in parentheses): vascular endothelial growth factor receptor (endothelium), albumin (liver), factor VII

(liver), fatty acid synthase (liver), von Willebrand factor (brain endothelium), alphaactin and myosin heavy chain (both in smooth muscle), synthetase I (small intestine), Na-K-Cl transporter (kidney). Additional cell type-specific TREs are known in the art.

5

AFP is an oncofetal protein, the expression of which is primarily restricted to developing tissues of endodermal origin (yolk sac, fetal liver, and gut), although the level of its expression varies greatly depending on the tissue and the developmental stage. AFP is of clinical interest because the serum concentration of AFP is elevated in a majority of hepatoma patients, with high levels of AFP found in patients with advanced disease. The serum AFP levels in patients appear to be regulated by AFP expression in hepatocellular carcinoma but not in surrounding normal liver. Thus, the AFP gene appears to be regulated to hepatoma cell-specific expression.

10

Cell type-specific TREs from the *AFP* gene have been identified. For example, the cloning and characterization of human AFP-specific enhancer activity is described in Watanabe et al. (1987) *J. Biol. Chem.* 262:4812-4818. The entire 5' *AFP* flanking region (containing the promoter, putative silencer, and enhancer elements) is contained within approximately 5 kb upstream from the transcription start site.

15

20

25

The *AFP* enhancer region in human is located between about nt -3954 and about nt -3335, relative to the transcription start site of the *AFP* gene. The human *AFP* promoter encompasses a region from about nt -174 to about nt +29. Juxtapositioning of these two genetic elements yields a fully functional *AFP*-TRE. Ido et al. (1995) describe a 259 bp promoter fragment (nt -230 to nt +29) that is specific for HCC. *Cancer Res.* 55:3105-3109. The *AFP* enhancer contains two regions, denoted A and B, located between nt -3954 and nt -3335 relative to the transcription start site. The promoter region contains typical TATA and CAAT boxes. Preferably, the *AFP*-TRE contains at least one enhancer region. More preferably, the *AFP*-TRE contains both enhancer regions.

Suitable target cells for adenoviral vectors containing AFP-TREs are any cell type that allow an AFP-TRE to function. Preferred are cells that express, or produce, AFP, including, but not limited to, tumor cells expressing AFP. Examples of such cells are hepatocellular carcinoma cells, gonadal and other germ cell tumors (especially endodermal sinus tumors), brain tumor cells, ovarian tumor cells, acinar cell carcinoma of the pancreas (Kawamoto et al. (1992) Hepatogastroenterology 39:282–286), primary gall bladder tumor (Katsuragi et al. (1989) Rinsko Hoshasen 34:371–374), uterine endometrial adenocarcinoma cells (Koyama et al. (1996) Jpn. J. Cancer Res. 87:612-617), and any metastases of the foregoing (which can occur in lung, adrenal gland, bone marrow, and/or spleen). In some cases, metastatic disease to the liver from certain pancreatic and stomach cancers produce AFP. Especially preferred are hepatocellular carcinoma cells and any of their metastases. AFP production can be measured using assays standard in the art, such as RIA, ELISA or Western blots (immunoassays) to determine levels of AFP protein production or Northern blots to determine levels of AFP mRNA production. Alternatively, such cells can be identified and/or characterized by their ability to activate transcriptionally an AFP-TRE (i.e., allow an AFP-TRE to function).

20

5

10

15

The protein urokinase plasminogen activator (uPA) and its cell surface receptor, urokinase plasminogen activator receptor (uPAR), are expressed in many of the most frequently occurring neoplasia and appear to represent important proteins in cancer metastasis. Both proteins are implicated in breast, colon, prostate, liver, renal, lung and ovarian cancer. Transcriptional regulatory elements that regulate uPA and uPAR transcription have been extensively studied. Riccio et al. (1985) *Nucleic Acids Res.* 13:2759-2771; Cannio et al. (1991) *Nucleic Acids Res.* 19:2303–2308.

25

CEA is a 180,000-Dalton glycoprotein tumor-associated antigen present on endodermally-derived neoplasia of the gastrointestinal tract, such as colorectal, gastric (stomach) and pancreatic cancer, as well as other adenocarcinomas such as breast and lung cancers. CEA is of clinical interest because circulating CEA can be

detected in the great majority of patients with CEA-positive tumors. In lung cancer, about 50% of total cases have circulating CEA, with high concentrations of CEA (greater than 20 ng/ml) often detected in adenocarcinomas. Approximately 50% of patients with gastric carcinoma are serologically positive for CEA.

The 5' upstream flanking sequence of the CEA gene has been shown to confer

5

cell-specific activity. The *CEA* promoter region, approximately the first 424 nucleotides upstream of the translational start site in the 5' flanking region of the gene, was shown to confer cell-specific activity when the region provided higher promoter activity in CEA-producing cells than in non-producing HeLa cells.. Schrewe et al. (1990) *Mol. Cell. Biol.* 10:2738-2748. In addition, cell-specific enhancer regions have been found. WO/95/14100. The entire 5' *CEA* flanking region (containing the promoter, putative silencer, and enhancer elements) appears to be contained within approximately 14.5 kb upstream from the transcription start site. Richards et al. (1995); WO 95/14100. Further characterization of the 5' flanking region of the *CEA* gene by Richards et al. (1995) indicated two upstream regions, -

15

10

13.6 to -10.7 kb or -6.1 to -4.0 kb, when linked to the multimerized promoter resulted in high-level and selective expression of a reporter construct in CEA-producing LoVo and SW1463 cells. Richards et al. (1995) also localized the promoter region to nt -90

20

25

essential for expression. WO95/14100 describes a series of 5' flanking *CEA* fragments which confer cell-specific activity, such as about nt -299 to about nt +69; about nt -90 to about nt +69; nt -14,500 to nt -10,600; nt -13,600 to nt -10,600, nt -

and nt +69 relative to the transcriptional start site, with region nt -41 to nt -18 as

.

6100 to nt -3800. In addition, cell specific transcription activity is conferred on an operably linked gene by the *CEA* fragment from nt -402 to nt +69, depicted in (SEQ

ID NO:6). Any *CEA*-TREs used in the present invention are derived from mammalian cells, including but not limited to, human cells. Thus, any of the *CEA*-TREs may be used in the invention as long as requisite desired functionality is displayed in the adenovirus vector. The cloning and characterization of *CEA* 

10

15

20

25

sequences have been described in the literature and are thus made available for practice of this invention and need not be described in detail herein.

The protein product of the *MUC1* gene (known as mucin or MUC1 protein; episialin; polymorphic epithelial mucin or PEM; EMA; DF3 antigen; NPGP; PAS-O; or CA15.3 antigen) is normally expressed mainly at the apical surface of epithelial cells lining the glands or ducts of the stomach, pancreas, lungs, trachea, kidney, uterus, salivary glands, and mammary glands. Zotter et al. (1988) *Cancer Rev.* 11–12: 55–101; and Girling et al. (1989) *Int. J. Cancer* 43: 1072–1076. However, mucin is overexpressed in 75–90% of human breast carcinomas. Kufe et al. (1984) *Hybridoma* 3: 223–232. For reviews, see Hilkens (1988) *Cancer Rev.* 11–12: 25–54; and Taylor-Papadimitriou, et al. (1990) *J. Nucl. Med. Allied Sci.* 34: 144–150. Mucin protein expression correlates with the degree of breast tumor differentiation. Lundy et al. (1985) *Breast Cancer Res. Treat.* 5: 269–276. This overexpression appears to be controlled at the transcriptional level.

Overexpression of the *MUC1* gene in human breast carcinoma cells MCF-7 and ZR-75-1 appears to be regulated at the transcriptional level. Kufe et al. (1984); Kovarik (1993) *J. Biol. Chem.* 268:9917–9926; and Abe et al. (1990) *J. Cell. Physiol.* 143: 226–231. The regulatory sequences of the *MUC1* gene have been cloned, including the approximately 0.9 kb upstream of the transcription start site which contains a TRE that appears to be involved in cell-specific transcription. Abe et al. (1993) *Proc. Natl. Acad. Sci. USA* 90: 282-286; Kovarik et al. (1993); and Kovarik et al. (1996) *J. Biol. Chem.* 271:18140–18147.

Any *MUC1*-TREs used in the present invention are derived from mammalian cells, including but not limited to, human cells. Preferably, the *MUC1*-TRE is human. In one embodiment, the *MUC1*-TRE may contain the entire 0.9 kb 5' flanking sequence of the *MUC1* gene. In other embodiments, the *MUC1*-TREs comprise the following sequences (relative to the transcription start site of the *MUC1* 

10

15

20

25

gene): about nt -725 to about nt +31, nt -743 to about nt +33, nt -750 to about nt +33, and nt -598 to about nt +485 (operably-linked to a promoter).

The c-erbB2/neu gene (HER-2/neu or HER) is a transforming gene that encodes a 185 kD epidermal growth factor receptor-related transmembrane glycoprotein. In humans, the c-erbB2/neu protein is expressed during fetal development, however, in adults, the protein is weakly detectable (by immunohistochemistry) in the epithelium of many normal tissues. Amplification and/or over-expression of the *c-erbB2/neu* gene has been associated with many human cancers, including breast, ovarian, uterine, prostate, stomach and lung cancers. The clinical consequences of the c-erbB2/neu protein over-expression have been best studied in breast and ovarian cancer. c-erbB2/neu protein over-expression occurs in 20 to 40% of intraductal carcinomas of the breast and 30% of ovarian cancers, and is associated with a poor prognosis in subcategories of both diseases. Human, rat and mouse c-erbB2/neu TREs have been identified and shown to confer c-erbB2/neu expressing cell specific activity. Tal et al. (1987) Mol. Cell. Biol. 7:2597–2601; Hudson et al. (1990) J. Biol. Chem. 265:4389-4393; Grooteclaes et al. (1994) Cancer Res. 54:4193–4199; Ishii et al. (1987) Proc. Natl. Acad. Sci. USA 84:4374–4378; Scott et al. (1994) J. Biol. Chem. 269:19848-19858.

The cell type-specific TREs listed above are provided as non-limiting examples of TREs that would function in the instant invention. Additional cell-specific TREs are known in the art, as are methods to identify and test cell specificity of suspected TREs.

#### Using the adenoviral vectors of the invention

The adenoviral vectors can be used in a variety of forms, including, but not limited to, naked polynucleotide (usually DNA) constructs; polynucleotide constructs complexed with agents to facilitate entry into cells, such as cationic liposomes or other cationic compounds such as polylysine; packaged into infectious adenovirus particles (which may render the adenoviral vector(s) more immunogenic); packaged

into other particulate viral forms such as HSV or AAV; complexed with agents (such as PEG) to enhance or dampen an immune response; complexed with agents that facilitate *in vivo* transfection, such as DOTMA<sup>TM</sup>, DOTAP<sup>TM</sup>, and polyamines. Thus, the invention also provides an adenovirus capable of replicating preferentially in cell status-producing cells. "Replicating preferentially" means that the adenovirus replicates more in cell exhibiting a requisite physiological state than a cell not exhibiting that state. Preferably, the adenovirus replicates at least about 2-fold higher, preferably at least about 5-fold higher, more preferably at least about 10-fold higher, still more preferably at least about 50-fold higher, even more preferably at least about 100-fold higher, still more preferably at least about 400-fold to about 500-fold higher, still more preferably at least about 1000-fold higher, most preferably at least about 1 x 10<sup>6</sup> higher. Most preferably, the adenovirus replicates solely in cells exhibiting a requisite physiological state (that is, does not replicate or replicates at very low levels in cells not exhibiting the requisite physiological state).

15

10

5

If an adenoviral vector is packaged into an adenovirus, the adenovirus itself may also be selected to further enhance targeting. For example, adenovirus fibers mediate primary contact with cellular receptor(s) aiding in tropism. See, e.g., Amberg et al. (1997) *Virol.* 227:239-244. If a particular subgenus of an adenovirus serotype displayed tropism for a target cell type and/or reduced affinity for non-target cell types, such subgenus (or subgenera) could be used to further increase cell-specificity of cytoxicity and/or cytolysis.

20

The adenoviral vectors may be delivered to the target cell in a variety of ways, including, but not limited to, liposomes, general transfection methods that are well known in the art (such as calcium phosphate precipitation or electroporation), direct injection, and intravenous infusion. The means of delivery will depend in large part on the particular adenoviral vector (including its form) as well as the type and location of the target cells (i.e., whether the cells are *in vitro* or *in vivo*).

If used as a packaged adenovirus, adenovirus vectors may be administered in an appropriate physiologically acceptable carrier at a dose of about 10<sup>4</sup> to about 10<sup>14</sup>. The multiplicity of infection will generally be in the range of about 0.001 to 100. If administered as a polynucleotide construct (i.e., not packaged as a virus) about 0.01 µg to about 1000 µg of an adenoviral vector can be administered. The adenoviral vector(s) may be administered one or more times, depending upon the intended use and the immune response potential of the host, and may also be administered as multiple, simultaneous injections. If an immune response is undesirable, the immune response may be diminished by employing a variety of immunosuppressants, so as to permit repetitive administration, without a strong immune response. If packaged as another viral form, such as HSV, an amount to be administered is based on standard knowledge about that particular virus (which is readily obtainable from, for example, published literature) and can be determined empirically.

15

20

10

5

#### Host cells comprising the adenoviral vectors of the invention

The present invention also provides host cells comprising (i.e., transformed with) the adenoviral vectors described herein. Both prokaryotic and eukaryotic host cells can be used as long as sequences requisite for maintenance in that host, such as appropriate replication origin(s), are present. For convenience, selectable markers are also provided. Prokaryotic host cells include bacterial cells, for example, *E. coli* and mycobacteria. Among eukaryotic host cells are yeast, insect, avian, plant and mammalian. Host systems are known in the art and need not be described in detail herein.

#### Compositions of the invention

25

The present invention also provides compositions, including pharmaceutical compositions, containing the adenoviral vectors described herein. Such compositions (especially pharmaceutical compositions) are useful for administration *in vivo*, for example, when measuring the degree of transduction and/or effectiveness of cell

killing in an individual. Pharmaceutical compositions, comprised an adenoviral vector of this invention in a pharmaceutically acceptable excipient (generally an effective amount of the adenoviral vector), are suitable for systemic administration to individuals in unit dosage forms, sterile parenteral solutions or suspensions, sterile non-parenteral solutions or oral solutions or suspensions, oil in water or water in oil emulsions and the like. Formulations for parenteral and nonparenteral drug delivery are known in the art and are set forth in *Remington's Pharmaceutical Sciences*, 19th Edition, Mack Publishing (1995). Pharmaceutical compositions also include lyophilized and/or reconstituted forms of the adenoviral vectors (including those packaged as a virus, such as adenovirus) of the invention.

Other compositions are used, and are useful for, detection methods described herein. For these compositions, the adenoviral vector usually is suspended in an appropriate solvent or solution, such as a buffer system. Such solvent systems are well known in the art.

15

20

5

10

#### Kits of the invention

The present invention also encompasses kits containing an adenoviral vector(s) of this invention. These kits can be used for diagnostic and/or monitoring purposes, preferably monitoring. Procedures using these kits can be performed by clinical laboratories, experimental laboratories, medical practitioners, or private individuals. Kits embodied by this invention allow someone to detect the presence of cell status-producing cells in a suitable biological sample, such as biopsy specimens.

The kits of the invention comprise an adenoviral vector described herein in suitable packaging. The kit may optionally provide additional components that are useful in the procedure, including, but not limited to, buffers, developing reagents, labels, reacting surfaces, means for detection, control samples, instructions, and interpretive information.

10

15

20

#### Preparation of the adenovirus vectors of the invention

The adenovirus vectors of this invention can be prepared using recombinant techniques that are standard in the art. Generally, a cell status-specific TRE is inserted 5' to the adenoviral gene of interest, preferably one or more early genes (although late gene(s) may be used). A cell status-specific TRE can be prepared using oligonucleotide synthesis (if the sequence is known) or recombinant methods (such as PCR and/or restriction enzymes). Convenient restriction sites, either in the natural adeno-DNA sequence or introduced by methods such as oligonucleotide directed mutagenesis and PCR, provide an insertion site for a cell status-specific TRE. Accordingly, convenient restriction sites for annealing (i.e., inserting) a cell status-specific TRE can be engineered onto the 5' and 3' ends of a cell status-specific TRE using standard recombinant methods, such as PCR.

Polynucleotides used for making adenoviral vectors of this invention may be obtained using standard methods in the art, such as chemical synthesis, by recombinant methods, and/or by obtaining the desired sequence(s) from biological sources.

Adenoviral vectors are conveniently prepared by employing two plasmids, one plasmid providing for the left hand region of adenovirus and the other plasmid providing for the right hand region, where the two plasmids share at least about 500 nt of middle region for homologous recombination. In this way, each plasmid, as desired, may be independently manipulated, followed by cotransfection in a competent host, providing complementing genes as appropriate, or the appropriate transcription factors for initiation of transcription from a cell status-specific TRE for propagation of the adenovirus. Plasmids are generally introduced into a suitable host cell such as 293 cells using appropriate means of transduction, such as cationic liposomes. Alternatively, *in vitro* ligation of the right and left-hand portions of the adenovirus genome can also be used to construct recombinant adenovirus derivative containing all the replication-essential portions of adenovirus genome. Berkner et al.

10

15

20

25

(1983) Nucleic Acid Research 11: 6003-6020; Bridge et al. (1989) J. Virol. 63: 631-638.

For convenience, plasmids are available that provide the necessary portions of adenovirus. Plasmid pXC.1 (McKinnon (1982) *Gene* 19:33-42) contains the wild-type left-hand end of Ad5. pBHG10 (Bett et al. (1994) *Proc. Natl. Acad. Sci USA* 91:8802-8806; Microbix Biosystems Inc., Toronto) provides the right-hand end of Ad5, with a deletion in E3. The deletion in E3 provides room in the virus to insert a 3 kb cell status-TRE without deleting the endogenous enhancer/promoter. Bett et al. (1994). The gene for E3 is located on the opposite strand from E4 (r-strand). pBHG11 provides an even larger E3 deletion (an additional 0.3 kb is deleted). Bett et al. (1994).

For manipulation of the early genes, the transcription start site of Ad5 E1A is at 498 and the ATG start site of the E1A protein is at 560 in the virus genome. This region can be used for insertion of an cell status-specific TRE. A restriction site may be introduced by employing polymerase chain reaction (PCR), where the primer that is employed may be limited to the Ad5 genome, or may involve a portion of the plasmid carrying the Ad5 genomic DNA. For example, where pBR322 is used, the primers may use the EcoRI site in the pBR322 backbone and the Xbal site at 1339 of Ad5. By carrying out the PCR in two steps, where overlapping primers at the center of the region introduce a 30 sequence change resulting in a unique restriction site, one can provide for insertion of heterologous TRE at that site.

A similar strategy may also be used for insertion of a heterologous TRE to regulate E1B. The E1B promoter of Ad5 consists of a single high-affinity recognition site for Spl and a TATA box. This region extends from 1636 to 1701. By insertion of a heterologous TRE in this region, one can provide for target cell-specific transcription of the E1B gene. By employing the left-hand region modified with a heterologous TRE regulating E1A as the template for introducing a heterologous TRE to regulate E1B, the resulting adenovirus vector will be dependent

upon the cell status-specific transcription factors for expression of both E1A and E1B.

Similarly, a cell status-specific TRE can be inserted upstream of the E2 gene to make its expression cell status specific. The E2 early promoter, mapping in Ad5 from 27050-27150, consists of a major and a minor transcription initiation site, the latter accounting for about 5% of the E2 transcripts, two non-canonical TATA boxes, two E2F transcription factor binding sites and an ATF transcription factor binding site. For a detailed review of the E2 promoter architecture see Swaminathan et al., *Curr. Topics in Micro. and Imm.* (1995) 199 (part 3):177-194.

10

15

5

For E4, one must use the right hand portion of the adenovirus genome. The E4 transcription start site is predominantly at 35609, the TATA box at 35638 and the first ATG/CTG of ORF 1 is at 35532. Virtanen et al. (1984) *J. Virol.* 51: 822-831. Using any of the above strategies for the other genes, a cell status-specific TRE may be introduced upstream from the transcription start site. For the construction of mutants in the E4 region, the co-transfection and homologous recombination are performed in W162 cells (Weinberg et al. (1983) *Proc. Natl. Acad. Sci.* 80:5383–5386) which provide E4 proteins *in trans* to complement defects in synthesis of these proteins. Alternatively, these constructs can be produced by *in vitro* ligation.

#### Methods using the adenovirus vectors of the invention

The adenoviral vectors of the invention can be used for a wide variety of purposes, which will vary with the desired or intended result. Accordingly, the present invention includes methods using the adenoviral vectors described above.

25

20

In one embodiment, methods are provided for conferring selective cytoxicity in target cells (i.e., cells exhibiting a requisite physiological state which allows a cell status-specific TRE to function), generally but not necessarily in an individual (preferably human), comprising contacting the cells with an adenovirus vector described herein, such that the adenovirus vector enters the cell. Cytotoxicity can be

10

15

20

measured using standard assays in the art, such as dye exclusion, <sup>3</sup>H-thymidine incorporation, and/or lysis.

In another embodiment, methods are provided for propagating an adenovirus specific for mammalian cells which allow a cell status-specific TRE to function. These methods entail combining an adenovirus vector with mammalian cells, whereby said adenovirus is propagated.

The invention further provides methods of suppressing tumor cell growth, generally but not necessarily in an individual (preferably human), comprising contacting a tumor cell with an adenoviral vector of the invention such that the adenoviral vector enters the tumor cell and exhibits selective cytotoxicity for the tumor cell. Tumor cell growth can be assessed by any means known in the art, including, but not limited to, measuring tumor size, determining whether tumor cells are proliferating using a <sup>3</sup>H-thymidine incorporation assay, or counting tumor cells.

The invention also includes methods for detecting target cells (i.e., cells which permit or induce a cell status-specific TRE to function) in a biological sample. These methods are particularly useful for monitoring the clinical and/or physiological condition of an individual (i.e., mammal), whether in an experimental or clinical setting. For these methods, cells of a biological sample are contacted with an adenovirus vector, and replication of the adenoviral vector is detected. A suitable biological sample is one in which cells exhibiting a requisite physiological (and/or environmental) state, for example, an aberrant physiological state (such as cells in hypoxic conditions and exhibiting a phenotype characteristic of cells in hypoxic conditions, such as expression of HIF-1) may be or are suspected to be present. Generally, in mammals, a suitable clinical sample is one in which cancerous cells exhibiting a requisite physiological state, such as cells within a solid tumor which are under hypoxic conditions, are suspected to be present. Such cells can be obtained, for example, by needle biopsy or other surgical procedure. Cells to be contacted may be treated to promote assay conditions, such as selective enrichment, and/or

solubilization. In these methods, target cells can be detected using *in vitro* assays that detect adenoviral proliferation, which are standard in the art. Examples of such standard assays include, but are not limited to, burst assays (which measure virus yield) and plaque assays (which measure infectious particles per cell). Propagation can also be detected by measuring specific adenoviral DNA replication, which are also standard assays.

The following examples are provided to illustrate but not limit the invention.

10

5

#### EXAMPLES

#### EXAMPLE 1

## Adenovirus vector comprising E1A under transcriptional control of a hypoxia responsive element and a PSA-TRE

General techniques

15

A human embryonic kidney cell line, 293, efficiently expresses E1A and E1B genes of Ad5 and exhibits a high transfection efficiency with adenovirus DNA. To generate recombinant adenovirus, 293 cells were co-transfected with one left end Ad5 plasmid and one right end Ad5 plasmid. Homologous recombination generates adenoviruses with the required genetic elements for replication in 293 cells which provide E1A and E1B proteins *in trans* to complement defects in synthesis of these proteins.

20

The plasmids to be combined were co-transfected into 293 cells using cationic liposomes such as Lipofectin (DOTMA:DOPE<sup>TM</sup>, Life Technologies) by combining the two plasmids, then mixing the plasmid DNA solution (10 μg of each plasmid in 500 μl of minimum essential medium (MEM) without serum or other additives) with a four-fold molar excess of liposomes in 200 μ1 of the same buffer. The DNA-lipid complexes were then placed on the cells and incubated at 37°C, 5% CO<sub>2</sub> for 16 hours. After incubation the medium was changed to MEM with 10% fetal bovine serum and

the cells are further incubated at 37°C, 5% CO<sub>2</sub>, for 10 days with two changes of medium. At the end of this time the cells and medium were transferred to tubes, freeze-thawed three times, and the lysate was used to infect 293 cells at the proper dilution to detect individual viruses as plaques.

5

Plaques obtained were plaque purified twice, and viruses were characterized for presence of desired sequences by PCR and occasionally by DNA sequencing. For further experimentation, the viruses were purified on a large scale by cesium chloride gradient centrifugation.

10

Adenovirus vectors in which E1A is under transcriptional control of a cell status-specific TRE

15

An adenovirus vector containing a hypoxia response element (HRE) was generated. CN796, an adenovirus vector in which E1A is under the control of a composite TRE consisting of an HRE and a PSA-TRE, was made by co-transfecting CN515 with pBHG10. CN515 was constructed by inserting a 67 base pair fragment from HRE eno1 (Jiang et al. (1997) *Cancer Research* 57:5328-5335) (SEQ ID NO:1; Figure 2) into CN65 at the BglII site. CN65 is a plasmid containing an enhancer and promoter from the human PSA gene, consisting of an enhancer from -5322 to -3738 fused to a PSA promoter from -541 to +12. This is the PSA-TRE contained within plasmid CN706. Rodriguez et al. (1997) *Cancer Res.* 57:2559-2563.

20

Virus growth in vitro

Growth selectivity of recombinant adenovirus is analyzed in plaque assays in which a single infectious particle produces a visible plaque by multiple rounds of infection and replication. Virus stocks are diluted to equal pfu/ml, then used to infect monolayers of cells for 1 hour. The inoculum is then removed and the cells are overlayed with semisolid agar containing medium and incubated at 37°C for 10 days. Plaques in the monolayer are then counted and titers of infectious virus on the various cells are calculated. The data are normalized to the titer of CN702 (wild type) on 293 cells.

10

15

#### Claims

#### What is claimed is:

- 1. An adenovirus vector comprising an adenovirus gene under transcriptional control of a transcriptional regulatory element (TRE) comprising a cell status-specific TRE.
- 2. The adenovirus vector of claim 1, wherein the adenovirus gene is essential for viral replication.
- 3. The adenovirus vector of claim 2, wherein the adenovirus gene is an early gene.
- 4. The adenovirus vector of claim 2, wherein the adenovirus gene is a late gene.
- 5. The adenovirus vector of claim 3, wherein the adenovirus early gene is E1A.
- 6. The adenovirus vector of claim 3, wherein the adenovirus early gene is E1B.
- 7. The adenovirus vector of claim 3, wherein the adenovirus early gene is E4.
- 8. The adenovirus vector of claim 1, wherein the cell status-specific TRE is human.
- 9. The adenovirus vector of claim 1, wherein the cell status-specific TRE comprises a hypoxia responsive element (HRE).
  - 10. The adenovirus vector of claim 9, wherein the HRE comprises SEQ ID NO:1.


15

- 11. The adenovirus vector of claim 1, wherein the cell status-specific TRE comprises a cell cycle specific element.
- 12. The adenovirus vector of claim 11, wherein the cell cycle-specific element is from the E2F-1 gene.
- 5 13. The adenovirus vector of claim 1, wherein the cell status-specific TRE comprises a heat-inducible element.
  - 14. The adenovirus vector of claim 1, further comprising a cell type-specific TRE.
  - 15. The adenovirus vector of claim 14, wherein the cell type-specific TRE is prostate cell specific.
  - 16. The adenovirus vector of claim 15, wherein the prostate cell-specific TRE is a *PSA*-TRE.
  - 17. The adenovirus vector of claim 1, further comprising a transgene under transcriptional control of a second cell status-specific TRE.
  - 18. An adenovirus vector comprising an adenovirus gene under transcriptional control of a a TRE comprising a cell status-specific TRE and a cell-type specific TRE.
  - 19 The adenovirus vector of claim 18, wherein the adenovirus gene is an early gene.
  - 20. The adenovirus vector of claim 19, wherein the adenovirus early gene is E1A.
  - 21. The adenovirus vector of claim 20, wherein the cell status-specific TRE comprises an HRE and the cell-type specific TRE is a PSA-TRE.

- 22. The adenovirus vector of claim 21, wherein the HRE comprises SEQ ID NO:1 and the PSA-TRE comprises nucleotides about 503 to about 2086 of SEQ ID NO:3 and nucleotides about 5285 to about 5836 of SEQ ID NO:3.
  - 23. A composition comprising an adenovirus vector of claim 1.
- 5 24. The composition of claim 23, further comprising a pharmaceutically acceptable excipient.
  - 25. A host cell comprising the adenovirus vector of claim 1.
  - 26. A method of propagating adenovirus specific for cells which allow a cell status-specific TRE to function, said method comprising combining an adenovirus according to claim 1 with the cells, whereby said adenovirus is propagated.
  - 27. A method for conferring selective cytotoxicity on a target cell, said method comprising contacting a cell which allows a cell status-specific TRE to function with an adenovirus vector of claim 1, whereby the vector enters the cell.
  - 28. A method for suppressing tumor growth comprising introducing the adenovirus vector of claim 1 into a tumor cell which allows a cell status-specific TRE to function, wherein introduction of the adenovirus vector results in suppression of tumor growth.

#### ABSTRACT OF THE DISCLOSURE

The present invention provides adenoviral vectors comprising cell status-specific transcriptional regulatory elements which confer cell status-specific transcriptional regulation on an adenoviral gene. A "cell status" is generally a reversible physiological and/or environmental state. The invention further provides compositions and host cells comprising the vectors, as well as methods of using the vectors.



ccccgagg cagtgcat gaggctcagg gcgtgcgt gagtcgcagcgagaccccg gggtgcag gccgga

ggcaggccgtg ggcagccaat cgcgtaaaag* cctcgccgag	655 655 655 655 655 655	ada cy ggcaa ggaag	ttttgta cgatcca acttcat tccttag gagtgag	ggccca cagaac tcctag tcctag tgaggca tgtgttt gtgcct
ggcgcgctaa tgtggcgggac gcctgccgcc agcgtcatg	agaagaga ggggcggg ggggcatt	totgggtgag tgaggatgga gggacagtct	atgggt cgccag attgta cacgaa atgaat	a c c p p c c c c b
agccaatagg ctcggcggct tttgcaggca atgggcccgc	ggggcc agtcag ggggcc	ata gtg	caccato attacao gtctgtt aggatgo ttcaago	$\omega$ $\alpha$ $\alpha$ $\alpha$ $\alpha$ $\alpha$ $\alpha$ $\alpha$
aaccgccgcc cgtggctctt gcggcggccg gccgccgccg	ttcccgtg cgcctggt	ccctcgc ggtgcag ggctggg	atcaggct ttccggtg tttctgtt tcagagca acagagca	tctgaagcca ggcagcctct agaccagtgt aaaaaggagtg gtaaatcctt gatctcgctc gacttctcgcc
gttgttcccg tcgcggcaaa ggggcggagc ccgcctgtca	atagaaag cccacgcc catccgga gccccatc	gattaggata gattaggata ctgtggaact	tatagaa gagaaaa aatgtat atataaa tggatgt	gtggcctaag ggcttagcct tgggaggagt tgtgtggcca gggagggctc tgtcgcccag gctcaagcaa atgcctggct
tcacggccgg aaggatttgg gggatcgagc cccgggccgc	cagtgggat cagcagggg gcccctcgc	atggg tcctg tcctg	gggctca ccagacg actgtga ttgttga attttgg	gaccaccctt ctgtttcttt cgctactgag gcattggagc cctagaagtg gctggtgtgc tcctcccacc
1321 1381 1441 1501	1081 1141 1201 1261	901 961 1021	54 661 721 781	61 1221 1241 1301 1421 1421 1421

## FIGURE 4A

aagcttctag	ttttctttc	ccggtgacat	cgtggaaagc	actagcatct	ctaagcaatg	60
atctgtgaca	atattcacag	tgtaatgcca	tccagggaac	tcaactgagc	cttgatgtcc	120
agagattttt	gtgtttttt	ctgagactga	gtctcgctct	gtgccaggct	ggagtgcagt	180
ggtgcaacct	tggctcactg	caagctccgc	ctcctgggtt	cacgccattc	tcctgcctca	240
gcctcctgag	tagctgggac	tacaggcacc	cgccaccacg	cctggctaat	ttttttgtat	300
ttttagtaga	gatggggttt	cactgtgtta	gccaggatgg	tctcagtctc	ctgacctcgt	360
gatctgccca	ccttggcctc	ccaaagtgct	gggatgacag	gcgtgagcca	ccgcgcctgg	420
ccgatatcca	gagattttt	ggggggctcc	atcacacaga	catgttgact	gtcttcatgg	480
ttgactttta	gtatccagcc	cctctagaaa	tctagctgat	atagtgtggc	tcaaaacctt	540
cagcacaaat	cacaccgtta	gactatctgg	tgtggcccaa	accttcaggt	gaacaaaggg	600
actctaatct	ggcaggatat	tccaaagcat	tagagatgac	ctcttgcaaa	gaaaaagaaa	660
tggaaaagaa	aaagaaagaa	aggaaaaaaa	aaaaaaaaa	gagatgacct	ctcaggctct	720
gaggggaaac	gcctgaggtc	tttgagcaag	gtcagtcctc	tgttgcacag	tctccctcac	780
agggtcattg	tgacgatcaa	atgtggtcac	gtgtatgagg	caccagcaca	tgcctggctc	840
tggggagtgc	cgtgtaagtg	tatgcttgca	ctgctgaatg	cttgggatgt	gtcagggatt	900
atcttcagca	cttacagatg	ctcatctcat	cctcacagca	tcactatggg	atgggtatta	960
ctggcctcat	ttgatggaga	aagtggctgt	ggctcagaaa	ggggggacca	ctagaccagg	1020
gacactctgg	atgctgggga	ctccagagac	catgaccact	caccaactgc	agagaaatta	1080
attgtggcct	gatgtccctg	tcctggagag	ggtggaggtg	gaccttcact	aacctcctac	1140
cttgaccctc	tcttttaggg	ctctttctga	cctccaccat	ggtactagga	ccccattgta	1200
ttctgtaccc	tcttgactct	atgaccccca	ctgcccactg	catccagctg	ggtcccctcc	1260
tatctctatt	cccagctggc	cagtgcagtc	tcagtgccca	cctgtttgtc	agtaactctg	1320
aaggggctga	cattttactg	acttgcaaac	aaataagcta	actttccaga	gttttgtgaa	1380
tgctggcaga	gtccatgaga	ctcctgagtc	agaggcaaag	gcttttactg	ctcacagctt	1440
agcagacagc	atgaggttca	tgttcacatt	agtacacctt	gccccccca	aatcttgtag	1500
ggtgaccaga	gcagtctagg	tggatgctgt	gcagaagggg	tttgtgccac	tggtgagaaa	1560
cctgagatta	ggaatcctca	atcttatact	gggacaactt	gcaaacctgc	tcagcctttg	1620
tctctgatga	agatattatc	ttcatgatct	tggattgaaa	acagacctac	tctggaggaa	1680

### FIGURE 4B

catattgtat	cgattgtcct	tgacagtaaa	caaatctgtt	gtaagagaca	ttatctttat	1740
tatctaggac	agtaagcaag	cctggatctg	agagagatat	catcttgcaa	ggatgcctgc	1800
tttacaaaca	tccttgaaac	aacaatccag	aaaaaaaag	gtgttgctgt	ctttgctcag	1860
aagacacaca	gatacgtgac	agaaccatgg	agaattgcct	cccaacgctg	ttcagccaga	1920
gccttccacc	cttgtctgca	ggacagtctc	aacgttccac	cattaaatac	ttcttctatc	1980
acatcctgct	tctttatgcc	taaccaaggt	tctaggtccc	gatcgactgt	gtctggcagc	2040
actccactgc	caaacccaga	ataaggcagc	gctcaggatc	ccgaaggggc	atggctgggg	2100
atcagaactt	ctgggtttga	gtgaggagtg	ggtccaccct	cttgaatttc	aaaggaggaa	2160
gaggctggat	gtgaaggtac	tgggggaggg	aaagtgtcag	ttccgaactc	ttaggtcaat	2220
gagggaggag	actggtaagg	tcccagctcc	cgaggtactg	atgtgggaat	ggcctaagaa	2280
tctcatatcc	tcaggaagaa	ggtgctggaa	tcctgagggg	tagagttctg	ggtatatttg	2340
tggcttaagg	ctctttggcc	cctgaaggca	gaggctggaa	ccattaggtc	cagggtttgg	2400
ggtgatagta	atgggatctc	ttgattcctc	aagagtctga	ggatcgaggg	ttgcccattc	2460
ttccatcttg	ccacctaatc	cttactccac	ttgagggtat	caccagccct	tctagctcca	2520
tgaaggtccc	ctgggcaagc	acaatctgag	catgaaagat	gccccagagg	ccttgggtgt	2580
catccactca	tcatccagca	tcacactctg	agggtgtggc	cagcaccatg	acgtcatgtt	2640
gctgtgacta	tccctgcagc	gtgcctctcc	agccacctgc	caaccgtaga	gctgcccatc	2700
ctcctctggt	gggagtggcc	tgcatggtgc	caggctgagg	cctagtgtca	gacagggagc	2760
ctggaatcat	agggatccag	gactcaaaag	tgctagagaa	tggccatatg	tcaccatcca	2820
tgaaatctca	agggcttctg	ggtggagggc	acagggacct	gaacttatgg	tttcccaagt	2880
ctattgctct	cccaagtgag	tctcccagat	acgaggcact	gtgccagcat	cagccttatc	2940
tccaccacat	cttgtaaaag	gactacccag	ggccctgatg	aacaccatgg	tgtgtacagg	3000
agtagggggt	ggaggcacgg	actcctgtga	ggtcacagcc	aagggagcat	catcatgggt	3060
ggggaggagg	caatggacag	gcttgagaac	ggggatgtgg	ttgtatttgg	ttttctttgg	3120
ttagataaag	tgctgggtat	aggattgaga	gtggagtatg	aagaccagtt	aggatggagg	3180
atcagattgg	agttgggtta	gataaagtgc	tgggtatagg	attgagagtg	gagtatgaag	3240
accagttagg	atggaggatc	agattggagt	tgggttagag	atggggtaaa	attgtgctcc	3300
ggatgagttt	gggattgaca	ctgtggaggt	ggtttgggat	ggcatggctt	tgggatggaa	3360

### FIGURE 4C

atagatttgt	tttgatgttg	gctcagacat	ccttggggat	tgaactgggg	atgaagctgg	3420
gtttgatttt	ggaggtagaa	gacgtggaag	tagctgtcag	atttgacagt	ggccatgagt	3480
tttgtttgat	ggggaatcaa	acaatggggg	aagacataag	ggttggcttg	ttaggttaag	3540
ttgcgttggg	ttgatggggt	cggggctgtg	tataatgcag	ttggattggt	ttgtattaaa	3600
ttgggttggg	tcaggttttg	gttgaggatg	agttgaggat	atgcttgggg	acaccggatc	3660
catgaggttc	tcactggagt	ggagacaaac	ttcctttcca	ggatgaatcc	agggaagcct	3720
taattcacgt	gtaggggagg	tcaggccact	ggctaagtat	atccttccac	tccagctcta	3780
agatggtctt	aaattgtgat	tatctatatc	cacttctgtc	tccctcactg	tgcttggagt	3840
ttacctgatc	actcaactag	aaacagggga	agattttatc	aaattctttt	ttttttttt	3900
tttttttga	gacagagtct	cactctgttg	cccaggctgg	agtgcagtgg	cgcagtctcg	3960
gctcactgca	acctctgcct	cccaggttca	agtgattctc	ctgcctcagc	ctcctgagtt	4020
gctgggatta	caggcatgca	gcaccatgcc	cagctaattt	ttgtattttt	agtagagatg	4080
gggtttcacc	aatgtttgcc	aggctggcct	cgaactcctg	acctggtgat	ccacctgcct	4140
cagcctccca	aagtgctggg	attacaggcg	tcagccaccg	cgcccagcca	cttttgtcaa	4200
attcttgaga	cacagctcgg	gctggatcaa	gtgagctact	ctggttttat	tgaacagctg	4260
aaataaccaa	ctttttggaa	attgatgaaa	tcttacggag	ttaacagtgg	aggtaccagg	4320
gctcttaaga	gttcccgatt	ctcttctgag	actacaaatt	gtgattttgc	atgccacctt	4380
aatcttttt	tttttttt	taaatcgagg	tttcagtctc	attctatttc	ccaggctgga	4440
gttcaatagc	gtgatcacag	ctcactgtag	ccttgaactc	ctggccttaa	gagattctcc	4500
tgcttcggtc	tcccaatagc	taagactaca	gtagtccacc	accatatcca	gataatttt	4560
aaattttttg	gggggccggg	cacagtggct	cacgcctgta	atcccaacac	catgggaggc	4620
tgagatgggt	ggatcacgag	gtcaggagtt	tgagaccagc	ctgaccaaca	tggtgaaact	4680
ctgtctctac	taaaaaaaaa	aaaaatagaa	aaattagccg	ggcgtggtgg	cacacggcac	4740
ctgtaatccc	agctactgag	gaggctgagg	caggagaatc	acttgaaccc	agaaggcaga	4800
ggttgcaatg	agccgagatt	gcgccactgc	actccagcct	gggtgacaga	gtgagactct	4860
gtctcaaaaa	aaaaaaattt	tttttttt	tttgtagaga	tggatcttgc	tttgtttctc	4920
tggttggcct	tgaactcctg	gcttcaagtg	atcctcctac	cttggcctcg	gaaagtgttg	4980
ggattacagg	cgtgagccac	catgactgac	ctgtcgttaa	tcttgaggta	cataaacctg	5040
gctcctaaag	gctaaaggct	aaatatttgt	tggagaaggg	gcattggatt	ttgcatgagg	5100

## FIGURE 4D

atgattctga	cctgggaggg	caggtcagca	ggcatctctg	ttgcacagat	agagtgtaca	5160
ggtctggaga	acaaggagtg	gggggttatt	ggaattccac	attgtttgct	gcacgttgga	5220
ttttgaaatg	ctagggaact	ttgggagact	catatttctg	ggctagagga	tctgtggacc	5280
acaagatctt	tttatgatga	cagtagcaat	gtatctgtgg	agctggattc	tgggttggga	5340
gtgcaaggaa	aagaatgtac	taaatgccaa	gacatctatt	tcaggagcat	gaggaataaa	5400
agttctagtt	tctggtctca	gagtggtgca	gggatcaggg	agtctcacaa	tctcctgagt	5460
gctggtgtct	tagggcacac	tgggtcttgg	agtgcaaagg	atctaggcac	gtgaggcttt	5520
gtatgaagaa	tcggggatcg	tacccacccc	ctgtttctgt	ttcatcctgg	gcatgtctcc	5580
tctgcctttg	tcccctagat	gaagtctcca	tgagctacaa	gggcctggtg	catccagggt	5640
gatctagtaa	ttgcagaaca	gcaagtgcta	gctctccctc	cccttccaca	gctctgggtg	5700
tgggaggggg	ttgtccagcc	tccagcagca	tggggagggc	cttggtcagc	ctctgggtgc	5760
cagcagggca	ggggcggagt	cctggggaat	gaaggtttta	tagggctcct	gggggaggct	5820
ccccagcccc	aagctt					5836

## FIGURE 5A

aagcttttta	gtgctttaga	cagtgagctg	gtctgtctaa	cccaagtgac	ctgggctcca	60
tactcagccc	cagaagtgaa	gggtgaagct	gggtggagcc	aaaccaggca	agcctaccct	120
cagggeteee	agtggcctga	gaaccattgg	acccaggacc	cattacttct	agggtaagga	180
aggtacaaac	accagatcca	accatggtct	ggggggacag	ctgtcaaatg	cctaaaaata	240
tacctgggag	aggagcaggc	aaactatcac	tgccccaggt	tctctgaaca	gaaacagagg	300
ggcaacccaa	agtccaaatc	caggtgagca	ggtgcaccaa	atgcccagag	atatgacgag	360
gcaagaagtg	aaggaaccac	ccctgcatca	aatgttttgc	atgggaagga	gaagggggtt	420
gctcatgttc	ccaatccagg	agaatgcatt	tgggatctgc	cttcttctca	ctccttggtt	480
agcaagacta	agcaaccagg	actctggatt	tggggaaaga	cgtttatttg	tggaggccag	540
tgatgacaat	cccacgaggg	cctaggtgaa	gagggcagga	aggctcgaga	cactggggac	600
tgagtgaaaa	ccacacccat	gatctgcacc	acccatggat	gctccttcat	tgctcacctt	660
tctgttgata	tcagatggcc	ccattttctg	taccttcaca	gaaggacaca	ggctagggtc	720
tgtgcatggc	cttcatcccc	ggggccatgt	gaggacagca	ggtgggaaag	atcatgggtc	780
ctcctgggtc	ctgcagggcc	agaacattca	tcacccatac	tgacctccta	gatgggaatg	840
gcttccctgg	ggctgggcca	acggggcctg	ggcaggggag	aaaggacgtc	aggggacagg	900
gaggaagggt	catcgagacc	cagcctggaa	ggttcttgtc	tctgaccatc	caggatttac	960
ttccctgcat	ctacctttgg	tcattttccc	tcagcaatga	ccagctctgc	ttcctgatct	1020
cagcctccca	ccctggacac	agcaccccag	tccctggccc	ggctgcatcc	acccaatacc	1080
ctgataaccc	aggacccatt	acttctaggg	taaggagggt	ccaggagaca	gaagctgagg	1140
aaaggtctga	agaagtcaca	tctgtcctgg	ccagagggga	aaaaccatca	gatgctgaac	1200
caggagaatg	ttgacccagg	aaagggaccg	aggacccaag	aaaggagtca	gaccaccagg	1260
gtttgcctga	gaggaaggat	caaggccccg	agggaaagca	gggctggctg	catgtgcagg	1320
acactggtgg	ggcatatgtg	tcttagattc	tccctgaatt	cagtgtccct	gccatggcca	1380
gactctctac	tcaggcctgg	acatgctgaa	ataggacaat	ggccttgtcc	tctctcccca	1440
ccatttggca	agagacataa	aggacattcc	aggacatgcc	ttcctgggag	gtccaggttc	1500
tctgtctcac	acctcaggga	ctgtagttac	tgcatcagcc	atggtaggtg	ctgatctcac	1560
ccagcctgtc	caggcccttc	cactctccac	tttgtgacca	tgtccaggac	cacccctcag	1620
atcctgagcc	tgcaaatacc	cccttgctgg	gtgggtggat	tcagtaaaca	gtgagctcct	1680

## FIGURE 5B

atccagcccc	cagagecace	tctgtcacct	tcctgctggg	catcatccca	ccttcacaag	1740
cactaaagag	catggggaga	cctggctagc	tgggtttctg	catcacaaag	aaaataatcc	1800
cccaggttcg	gattcccagg	gctctgtatg	tggagctgac	agacctgagg	ccaggagata	1860
gcagaggtca	gccctaggga	gggtgggtca	tccacccagg	ggacaggggt	gcaccagcct	1920
tgctactgaa	agggcctccc	caggacagcg	ccatcagccc	tgcctgagag	ctttgctaaa	1980
cagcagtcag	aggaggccat	ggcagtggct	gagctcctgc	tccaggcccc	aacagaccag	2040
accaacagca	caatgcagtc	cttccccaac	gtcacaggtc	accaaaggga	aactgaggtg	2100
ctacctaacc	ttagagccat	caggggagat	aacagcccaa	tttcccaaac	aggccagttt	2160
caatcccatg	acaatgacct	ctctgctctc	attcttccca	aaataggacg	ctgattctcc	2220
cccaccatgg	atttctccct	tgtcccggga	gccttttctg	ccccctatga	tctgggcact	2280
cctgacacac	acctcctctc	tggtgacata	tcagggtccc	tcactgtcaa	gcagtccaga	2340
aaggacagaa	ccttggacag	cgcccatctc	agcttcaccc	ttcctccttc	acagggttca	2400
gggcaaagaa	taaatggcag	aggccagtga	gcccagagat	ggtgacaggc	agtgacccag	2460
gggcagatgc	ctggagcagg	agctggcggg	gccacaggga	gaaggtgatg	caggaaggga	2520
aacccagaaa	tgggcaggaa	aggaggacac	aggctctgtg	gggctgcagc	ccagggttgg	2580
actatgagtg	tgaagccatc	tcagcaagta	aggccaggtc	ccatgaacaa	gagtgggagc	2640
acgtggcttc	ctgctctgta	tatggggtgg	gggattccat	gccccataga	accagatggc	2700
cggggttcag	atggagaagg	agcaggacag	gggatcccca	ggataggagg	accccagtgt	2760
ccccacccag	gcaggtgact	gatgaatggg	catgcagggt	cctcctgggc	tgggctctcc	2820
ctttgtccct	caggattcct	tgaaggaaca	tccggaagcc	gaccacatct	acctggtggg	2880
ttctggggag	tccatgtaaa	gccaggagct	tgtgttgcta	ggaggggtca	tggcatgtgc	2940
tgggggcacc	aaagagagaa	acctgagggc	aggcaggacc	tggtctgagg	aggcatggga	3000
gcccagatgg	ggagatggat	gtcaggaaag	gctgccccat	cagggagggt	gatagcaatg	3060
gggggtctgt	gggagtgggc	acgtgggatt	ccctgggctc	tgccaagttc	cctcccatag	3120
tcacaacctg	gggacactgc	ccatgaaggg	gcgcctttgc	ccagccagat	gctgctggtt	3180
ctgcccatcc	actaccctct	ctgctccagc	cactctgggt	ctttctccag	atgccctgga	3240
cagecetgge	ctgggcctgt	cccctgagag	gtgttgggag	aagctgagtc	tctggggaca	3300
ctctcatcag	agtctgaaag	gcacatcagg	aaacatccct	ggtctccagg	actaggcaat	3360

### FIGURE 5C

gaggaaaggg	ccccagctcc	tccctttgcc	actgagaggg	tcgaccctgg	gtggccacag	3420
tgacttctgc	gtctgtccca	gtcaccctga	aaccacaaca	aaaccccagc	cccagaccct	3480
gcaggtacaa	tacatgtggg	gacagtctgt	acccagggga	agccagttct	ctcttcctag	3540
gagaccgggc	ctcagggctg	tgcccggggc	aggcgggggc	agcacgtgcc	tgtccttgag	3600
aactcgggac	cttaagggtc	tctgctctgt	gaggcacagc	aaggatcctt	ctgtccagag	3660
atgaaagcag	ctcctgcccc	tcctctgacc	tcttcctcct	tcccaaatct	caaccaacaa	3720
ataggtgttt	caaatctcat	catcaaatct	tcatccatcc	acatgagaaa	gcttaaaacc	3780
caatggattg	acaacatcaa	gagttggaac	aagtggacat	ggagatgtta	cttgtggaaa	3840
tttagatgtg	ttcagctatc	gggcaggaga	atctgtgtca	aattccagca	tggttcagaa	3900
gaatcaaaaa	gtgtcacagt	ccaaatgtgc	aacagtgcag	gggataaaac	tgtggtgcat	3960
tcaaactgag	ggatattttg	gaacatgaga	aaggaaggga	ttgctgctgc	acagaacatg	4020
gatgatctca	cacatagagt	tgaaagaaag	gagtcaatcg	cagaatagaa	aatgatcact	4080
aattccacct	ctataaagtt	tccaagagga	aaacccaatt	ctgctgctag	agatcagaat	4140
ggaggtgacc	tgtgccttgc	aatggctgtg	agggtcacgg	gagtgtcact	tagtgcaggc	4200
aatgtgccgt	atcttaatct	gggcagggct	ttcatgagca	cataggaatg	cagacattac	4260
tgctgtgttc	attttacttc	accggaaaag	aagaataaaa	tcagccgggc	gcggtggctc	4320
acgcctgtaa	tcccagcact	ttagaaggct	gaggtgggca	gattacttga	ggtcaggagt	4380
tcaagaccac	cctggccaat	atggtgaaac	cccggctcta	ctaaaaatac	aaaaattagc	4440
tgggcatggt	ggtgcgcgcc	tgtaatccca	gctactcggg	aggctgaggc	tggacaattg	4500
cttggaccca	ggaagcagag	gttgcagtga	gccaagattg	tgccactgca	ctccagcttg	4560
ggcaacagag	ccagactctg	taaaaaaaaa	aaaaaaaaa	aaaaaaagaa	agaaagaaaa	4620
agaaaagaaa	gtataaaatc	tctttgggtt	aacaaaaaa	gatccacaaa	acaaacacca	4680
gctcttatca	aacttacaca	actctgccag	agaacaggaa	acacaaatac	tcattaactc	4740
acttttgtgg	caataaaacc	ttcatgtcaa	aaggagacca	ggacacaatg	aggaagtaaa	4800
actgcaggcc	ctacttgggt	gcagagaggg	aaaatccaca	aataaaacat	taccagaagg	4860
agctaagatt	tactgcattg	agttcattcc	ccaggtatgc	aaggtgattt	taacacctga	4920
aaatcaatca	ttgcctttac	tacatagaca	gattagctag	aaaaaatta	caactagcag	4980
aacagaagca	atttggcctt	cctaaaattc	cacatcatat	catcatgatg	gagacagtgc	5040
agacgccaat	gacaataaaa	agagggacct	ccgtcacccg	gtaaacatgt	ccacacagct	5100

#### FIGURE 5D

ccagcaagca	cccgtcttcc	cagtgaatca	ctgtaacctc	ccctttaatc	agccccaggc	5160
aaggctgcct	gcgatggcca	cacaggctcc	aacccgtggg	cctcaacctc	ccgcagaggc	5220
tctcctttgg	ccaccccatg	gggagagcat	gaggacaggg	cagagccctc	tgatgcccac	5280
acatggcagg	agctgacgcc	agagccatgg	gggctggaga	gcagagctgc	tggggtcaga	5340
gcttcctgag	gacacccagg	cctaagggaa	ggcagctccc	tggatggggg	caaccaggct	5400
ccgggctcca	acctcagagc	ccgcatggga	ggagccagca	ctctaggcct	ttcctagggt	5460
gactctgagg	ggaccctgac	acgacaggat	cgctgaatgc	acccgagatg	aaggggccac	5520
cacgggaccc	tgctctcgtg	gcagatcagg	agagagtggg	acaccatgcc	aggcccccat	5580
ggcatggctg	cgactgaccc	aggccactcc	cctgcatgca	tcagcctcgg	taagtcacat	5640
gaccaagccc	aggaccaatg	tggaaggaag	gaaacagcat	cccctttagt	gatggaaccc	5700
aaggtcagtg	caaagagagg	ccatgagcag	ttaggaaggg	tggtccaacc	tacagcacaa	5760
accatcatct	atcataagta	gaagccctgc	tccatgaccc	ctgcatttaa	ataaacgttt	5820
gttaaatgag	tcaaattccc	tcaccatgag	agctcacctg	tgtgtaggcc	catcacacac	5880
acaaacacac	acacacacac	acacacacac	acacacacac	acagggaaag	tgcaggatcc	5940
tggacagcac	caggcaggct	tcacaggcag	agcaaacagc	gtgaatgacc	catgcagtgc	6000
cctgggcccc	atcagctcag	agaccctgtg	agggctgaga	tggggctagg	caggggagag	6060
acttagagag	ggtggggcct	ccagggaggg	ggctgcaggg	agctgggtac	tgccctccag	6120
ggaggggct	gcagggagct	gggtactgcc	ctccagggag	ggggctgcag	ggagctgggt	6180
actgccctcc	agggagggg	ctgcagggag	ctgggtactg	ccctccaggg	agggggctgc	6240
agggagctgg	gtactgccct	ccagggaggc	aggagcactg	ttcccaacag	agagcacatc	6300
ttcctgcagc	agctgcacag	acacaggagc	ccccatgact	gccctgggcc	agggtgtgga	6360
ttccaaattt	cgtgccccat	tgggtgggac	ggaggttgac	cgtgacatcc	aaggggcatc	6420
tgtgattcca	aacttaaact	actgtgccta	caaaatagga	aataacccta	ctttttctac	6480
tatctcaaat	tccctaagca	caagctagca	ccctttaaat	caggaagttc	agtcactcct	6540
ggggtcctcc	catgccccca	gtctgacttg	caggtgcaca	gggtggctga	catctgtcct	6600
tgctcctcct	cttggctcaa	ctgccgcccc	tcctgggggt	gactgatggt	caggacaagg	6660
gatcctagag	ctggccccat	gattgacagg	aaggcaggac	ttggcctcca	ttctgaagac	6720
taggggtgtc	aagagagctg	ggcatcccac	agagctgcac	aagatgacgc	ggacagaggg	6780

#### FIGURE 5E

tgacacaggg	ctcagggctt	cagacgggtc	gggaggctca	gctgagagtt	cagggacaga	6840
cctgaggagc	ctcagtggga	aaagaagcac	tgaagtggga	agttctggaa	tgttctggac	6900
aagcctgagt	gctctaagga	aatgctccca	ccccgatgta	gcctgcagca	ctggacggtc	6960
tgtgtacctc	cccgctgccc	atcctctcac	agcccccgcc	tctagggaca	caactcctgc	7020
cctaacatgc	atctttcctg	tctcattcca	cacaaaaggg	cctctggggt	ccctgttctg	7080
cattgcaagg	agtggaggtc	acgttcccac	agaccaccca	gcaacagggt	cctatggagg	7140
tgcggtcagg	aggatcacac	gtccccccat	gcccagggga	ctgactctgg	gggtgatgga	7200
ttggcctgga	ggccactggt	cccctctgtc	cctgagggga	atctgcaccc	tggaggctgc	7260
cacatccctc	ctgattcttt	cagctgaggg	cccttcttga	aatcccaggg	aggactcaac	7320
ccccactggg	aaaggcccag	tgtggacggt	tccacagcag	cccagctaag	gcccttggac	7380
acagatcctg	agtgagagaa	cctttaggga	cacaggtgca	cggccatgtc	cccagtgccc	7440
acacagagca	ggggcatctg	gaccctgagt	gtgtagctcc	cgcgactgaa	cccagccctt	7500
ccccaatgac	gtgacccctg	gggtggctcc	aggtctccag	tccatgccac	caaaatctcc	7560
agattgaggg	tectecettg	agtccctgat	gcctgtccag	gagctgcccc	ctgagcaaat	7620
ctagagtgca	gagggctggg	attgtggcag	taaaagcagc	cacatttgtc	tcaggaagga	7680
aagggaggac	atgagctcca	ggaagggcga	tggcgtcctc	tagtgggcgc	ctcctgttaa	7740
tgagcaaaaa	ggggccagga	gagttgagag	atcagggctg	gccttggact	aaggctcaga	7800
tggagaggac	tgaggtgcaa	agagggggct	gaagtagggg	agtggtcggg	agagatggga	7860
ggagcaggta	aggggaagcc	ccagggaggc	cgggggaggg	tacagcagag	ctctccactc	7920
ctcagcattg	acatttgggg	tggtcgtgct	agtggggttc	tgtaagttgt	agggtgttca	7980
gcaccatctg	gggactctac	ccactaaatg	ccagcaggac	tccctcccca	agctctaaca	8040
accaacaatg	tctccagact	ttccaaatgt	cccctggaga	gcaaaattgc	ttctggcaga	8100
atcactgatc	tacgtcagtc	tctaaaagtg	actcatcagc	gaaatccttc	acctcttggg	8160
agaagaatca	caagtgtgag	aggggtagaa	actgcagact	tcaaaatctt	tccaaaagag	8220
ttttacttaa	tcagcagttt	gatgtcccag	gagaagatac	atttagagtg	tttagagttg	8280
atgccacatg	gctgcctgta	cctcacagca	ggagcagagt	gggttttcca	agggcctgta	8340
accacaactg	gaatgacact	cactgggtta	cattacaaag	tggaatgtgg	ggaattctgt	8400
agactttggg	aagggaaatg	tatgacgtga	gcccacagcc	taaggcagtg	gacagtccac	8460
tttgaggctc	tcaccatcta	ggagacatct	cagccatgaa	catagccaca	tctgtcatta	8520

## FIGURE 5F

gaaaacatgt	tttattaaga	ggaaaaatct	aggctagaag	tgctttatgc	tctttttct	8580
ctttatgttc	aaattcatat	acttttagat	cattccttaa	agaagaatct	atccccctaa	8640
gtaaatgtta	tcactgactg	gatagtgttg	gtgtctcact	cccaacccct	gtgtggtgac	8700
agtgccctgc	ttccccagcc	ctgggccctc	tctgattcct	gagagctttg	ggtgctcctt	8760
cattaggagg	aagagaggaa	gggtgttttt	aatattctca	ccattcaccc	atccacctct	8820
tagacactgg	gaagaatcag	ttgcccactc	ttggatttga	tcctcgaatt	aatgacctct	8880
atttctgtcc	cttgtccatt	tcaacaatgt	gacaggccta	agaggtgcct	tctccatgtg	8940
atttttgagg	agaaggttct	caagataagt	tttctcacac	ctctttgaat	tacctccacc	9000
tgtgtcccca	tcaccattac	cagcagcatt	tggacccttt	ttctgttagt	cagatgcttt	9060
ccacctcttg	agggtgtata	ctgtatgctc	tctacacagg	aatatgcaga	ggaaatagaa	9120
aaagggaaat	cgcattacta	ttcagagaga	agaagacctt	tatgtgaatg	aatgagagtc	9180
taaaatccta	agagagccca	tataaaatta	ttaccagtgc	taaaactaca	aaagttacac	9240
taacagtaaa	ctagaataat	aaaacatgca	tcacagttgc	tggtaaagct	aaatcagata	9300
tttttttctt	agaaaaagca	ttccatgtgt	gttgcagtga	tgacaggagt	gcccttcagt	9360
caatatgctg	cctgtaattt	ttgttccctg	gcagaatgta	ttgtcttttc	tccctttaaa	9420
tcttaaatgc	aaaactaaag	gcagctcctg	ggccccctcc	ccaaagtcag	ctgcctgcaa	9480
ccagccccac	gaagagcaga	ggcctgagct	tccctggtca	aaataggggg	ctagggagct	9540
taaccttgct	cgataaagct	gtgttcccag	aatgtcgctc	ctgttcccag	gggcaccagc	9600
ctggagggtg	gtgagcctca	ctggtggcct	gatgcttacc	ttgtgccctc	acaccagtgg	9660
tcactggaac	cttgaacact	tggctgtcgc	ccggatctgc	agatgtcaag	aacttctgga	9720
agtcaaatta	ctgcccactt	ctccagggca	gatacctgtg	aacatccaaa	accatgccac	9780
agaaccctgc	ctggggtcta	caacacatat	ggactgtgag	caccaagtcc	agccctgaat	9840
ctgtgaccac	ctgccaagat	gcccctaact	gggatccacc	aatcactgca	catggcaggc	9900
agcgaggctt	ggaggtgctt	cgccacaagg	cagccccaat	ttgctgggag	tttcttggca	9960
cctggtagtg	gtgaggagcc	ttgggaccct	caggattact	ccccttaagc	atagtgggga	10020
cccttctgca	tccccagcag	gtgccccgct	cttcagagcc	tctctctctg	aggtttaccc	10080
agacccctgc	accaatgaga	ccatgctgaa	gcctcagaga	gagagatgga	gctttgacca	10140
ggagccgctc	ttccttgagg	gccagggcag	ggaaagcagg	aggcagcacc	aggagtggga	10200

#### FIGURE 5G

acaccagtgt	ctaagcccct	gatgagaaca	gggtggtctc	tcccatatgc	ccataccagg	10260
cctgtgaaca	gaatcctcct	tctgcagtga	caatgtctga	gaggacgaca	tgtttcccag	10320
cctaacgtgc	agccatgccc	atctacccac	tgcctactgc	aggacagcac	caacccagga	10380
gctgggaagc	tgggagaaga	catggaatac	ccatggcttc	tcaccttcct	ccagtccagt	10440
gggcaccatt	tatgcctagg	acacccacct	gccggcccca	ggctcttaag	agttaggtca	10500
cctaggtgcc	tctgggaggc	cgaggcagga	gaattgcttg	aacccgggag	gcagaggttg	10560
cagtgagccg	agatcacacc	actgcactcc	agcctgggtg	acagaatgag	actctgtctc	10620
aaaaaaaag	agaaagatag	catcagtggc	taccaagggc	taggggcagg	ggaaggtgga	10680
gagttaatga	ttaatagtat	gaagtttcta	tgtgagatga	tgaaaatgtt	ctggaaaaaa	10740
aaatatagtg	gtgaggatgt	agaatattgt	gaatataatt	aacggcattt	aattgtacac	10800
ttaacatgat	taatgtggca	tattttatct	tatgtatttg	actacatcca	agaaacactg	10860
ggagagggaa	agcccaccat	gtaaaataca	cccaccctaa	tcagatagtc	ctcattgtac	10920
ccaggtacag	gcccctcatg	acctgcacag	gaataactaa	ggatttaagg	acatgaggct	10980
tcccagccaa	ctgcaggtgc	acaacataaa	tgtatctgca	aacagactga	gagtaaagct	11040
gggggcacaa	acctcagcac	tgccaggaca	cacacccttc	tcgtggattc	tgactttatc	11100
tgacccggcc	cactgtccag	atcttgttgt	gggattggga	caagggaggt	cataaagcct	11160
gtccccaggg	cactctgtgt	gagcacacga	gacctcccca	ccccccacc	gttaggtctc	11220
cacacataga	tctgaccatt	aggcattgtg	aggaggactc	tagcgcgggc	tcagggatca	11280
caccagagaa	tcaggtacag	agaggaagac	ggggctcgag	gagctgatgg	atgacacaga	11340
gcagggttcc	tgcagtccac	aggtccagct	caccctggtg	taggtgcccc	atccccctga	11400
tccaggcatc	cctgacacag	ctccctcccg	gagcctcctc	ccaggtgaca	catcagggtc	11460
cctcactcaa	gctgtccaga	gagggcagca	ccttggacag	cgcccacccc	acttcactct	11520
tectecetea	cagggctcag	ggctcagggc	tcaagtctca	gaacaaatgg	cagaggccag	11580
tgagcccaga	gatggtgaca	gggcaatgat	ccaggggcag	ctgcctgaaa	cgggagcagg	11640
tgaagccaca	gatgggagaa	gatggttcag	gaagaaaaat	ccaggaatgg	gcaggagagg	11700
agaggaggac	acaggctctg	tggggctgca	gcccaggatg	ggactaagtg	tgaagacatc	11760
tcagcaggtg	aggccaggtc	ccatgaacag	agaagcagct	cccacctccc	ctgatgcacg	11820
gacacacaga	gtgtgtggtg	ctgtgccccc	agagtcgggc	tctcctgttc	tggtccccag	11880
ggagtgagaa	gtgaggttga	cttgtccctg	ctcctctctg	ctaccccaac	attcaccttc	11940

### FIGURE 5H

tcctcatgcc	cctctctctc	aaatatgatt	tggatctatg	tccccgccca	aatctcatgt	12000
caaattgtaa	accccaatgt	tggaggtggg	gccttgtgag	aagtgattgg	ataatgcggg	12060
tggattttct	gctttgatgc	tgtttctgtg	atagagatct	cacatgatct	ggttgtttaa	12120
aagtgtgtag	cacctctccc	ctctctctct	ctctctctta	ctcatgctct	gccatgtaag	12180
acgttcctgt	ttccccttca	ccgtccagaa	tgattgtaag	ttttctgagg	cctccccagg	12240
agcagaagcc	actatgcttc	ctgtacaact	gcagaatgat	gagcgaatta	aacctctttt	12300
ctttataaat	tacccagtct	caggtatttc	tttatagcaa	tgcgaggaca	gactaataca	12360
atcttctact	cccagatccc	cgcacacgct	tagccccaga	catcactgcc	cctgggagca	12420
tgcacagcgc	agcctcctgc	cgacaaaagc	aaagtcacaa	aaggtgacaa	aaatctgcat	12480
ttggggacat	ctgattgtga	aagagggagg	acagtacact	tgtagccaca	gagactgggg	12540
ctcaccgagc	tgaaacctgg	tagcactttg	gcataacatg	tgcatgaccc	gtgttcaatg	12600
tctagagatc	agtgttgagt	aaaacagcct	ggtctggggc	cgctgctgtc	cccacttccc	12660
tcctgtccac	cagagggcgg	cagagttcct	cccaccctgg	agcctcccca	ggggctgctg	12720
acctccctca	gccgggccca	cagcccagca	gggtccaccc	tcacccgggt	cacctcggcc	12780
cacgtcctcc	tegeceteeg	agctcctcac	acggactctg	tcagctcctc	cctgcagcct	12840
atcggccgcc	cacctgaggc	ttgtcggccg	cccacttgag	gcctgtcggc	.tgccctctgc	12900
aggcagctcc	tgtcccctac	accccctcct	tccccgggct	cagctgaaag	ggcgtctccc	12960
agggcagctc	cctgtgatct	ccaggacagc	tcagtctctc	acaggeteeg	acgcccccta	13020
tgctgtcacc	tcacagccct	gtcattacca	ttaactcctc	agtcccatga	agttcactga	13080
gcgcctgtct	cccggttaca	ggaaaactct	gtgacaggga	ccacgtctgt	cctgctctct	13140
gtggaatccc	agggcccagc	ccagtgcctg	acacggaaca	gatgctccat	aaatactggt	13200
taaatgtgtg	ggagatctct	aaaaagaagc	atatcacctc	cgtgtggccc	ccagcagtca	13260
gagtctgttc	catgtggaca	caggggcact	ggcaccagca	tgggaggagg	ccagcaagtg	13320
cccgcggctg	ccccaggaat	gaggcctcaa	ccccagagc	ttcagaaggg	aggacagagg	13380
cctgcaggga	atagatcctc	cggcctgacc	ctgcagccta	atccagagtt	cagggtcagc	13440
tcacaccacg	tcgaccctgg	tcagcatccc	tagggcagtt	ccagacaagg	ccggaggtct	13500
cctcttgccc	tccagggggt	gacattgcac	acagacatca	ctcaggaaac	ggattcccct	13560
ggacaggaac	ctggctttgc	taaggaagtg	gaggtggagc	ctggtttcca	tcccttgctc	13620

#### FIGURE 5I

caacagaccc	ttctgatctc	tcccacatac	ctgctctgtt	cctttctggg	tcctatgagg	13680
accctgttct	gccaggggtc	cctgtgcaac	tccagactcc	ctcctggtac	caccatgggg	13740
aaggtggggt	gatcacagga	cagtcagcct	cgcagagaca	gagaccaccc	aggactgtca	13800
gggagaacat	ggacaggccc	tgagccgcag	ctcagccaac	agacacggag	agggagggtc	13860
cccctggagc	cttccccaag	gacagcagag	cccagagtca	cccacctccc	tccaccacag	13920
tcctctcttt	ccaggacaca	caagacacct	cccctccac	atgcaggatc	tggggactcc	13980
tgagacctct	gggcctgggt	ctccatccct	gggtcagtgg	cggggttggt	ggtactggag	14040
acagagggct	ggtccctccc	cagccaccac	ccagtgagcc	tttttctagc	ccccagagcc	14100
acctctgtca	ccttcctgtt	gggcatcatc	ccaccttccc	agagccctgg	agagcatggg	14160
gagacccggg	accctgctgg	gtttctctgt	cacaaaggaa	aataatcccc	ctggtgtgac	14220
agacccaagg	acagaacaca	gcagaggtca	gcactgggga	agacaggttg	tcctcccagg	14280
ggatgggggt	ccatccacct	tgccgaaaag	atttgtctga	ggaactgaaa	atagaaggga	14340
aaaaagagga	gggacaaaag	aggcagaaat	gagagggag	gggacagagg	acacctgaat	14400
aaagaccaca	cccatgaccc	acgtgatgct	gagaagtact	cctgccctag	gaagagactc	14460
agggcagagg	gaggaaggac	agcagaccag	acagtcacag	cagccttgac	aaaacgttcc	14520
tggaactcaa	gctcttctcc	acagaggagg	acagagcaga	cagcagagac	catggagtct	14580
ccctcggccc	ctccccacag	atggtgcatc	ccctggcaga	ggctcctgct	cacaggtgaa	14640
gggaggacaa	cctgggagag	ggtgggagga	gggagctggg	gtctcctggg	taggacaggg	14700
ctgtgagacg	gacagagggc	tcctgttgga	gcctgaatag	ggaagaggac	atcagagagg	14760
gacaggagtc	acaccagaaa	aatcaaattg	aactggaatt	ggaaaggggc	aggaaaacct	14820
caagagttct	attttcctag	ttaattgtca	ctggccacta	cgtttttaaa	aatcataata	14880
actgcatcag	atgacacttt	aaataaaaac	ataaccaggg	catgaaacac	tgtcctcatc	14940
cgcctaccgc	ggacattgga	aaataagccc	caggctgtgg	agggccctgg	gaaccctcat	15000
gaactcatcc	acaggaatct	gcagcctgtc	ccaggcactg	gggtgcaacc	aagatc	15056

### FIGURE 6A

gaattcagaa	ataggggaag	gttgaggaag	gacactgaac	tcaaagggga	tacagtgatt	60
ggtttatttg	tcttctcttc	acaacattgg	tgctggagga	attcccaccc	tgaggttatg	120
aagatgtctg	aacacccaac	acatagcact	ggagatatga	gctcgacaag	agtttctcag	180
ccacagagat	tcacagccta	gggcaggagg	acactgtacg	ccaggcagaa	tgacatggga	240
attgcgctca	cgattggctt	gaagaagcaa	ggactgtggg	aggtgggctt	tgtagtaaca	300
agagggcagg	gtgaactctg	attcccatgg	gggaatgtga	tggtcctgtt	acaaattttt	360
caagctggca	gggaataaaa	cccattacgg	tgaggacctg	tggagggcgg	ctgccccaac	420
tgataaagga	aatagccagg	tgggggcctt	tcccattgta	ggggggacat	atctggcaat	480
agaagccttt	gagacccttt	agggtacaag	tactgaggca	gcaaataaaa	tgaaatctta	540
tttttcaact	ttatactgca	tgggtgtgaa	gatatatttg	tttctgtaca	gggggtgagg	600
gaaaggaggg	gaggaggaaa	gttcctgcag	gtctggtttg	gtcttgtgat	ccagggggtc	660
ttggaactat	ttaaattaaa	ttaaattaaa	acaagcgact	gttttaaatt	aaattaaatt	720
aaattaaatt	ttactttatt	ttatcttaag	ttctgggcta	catgtgcagg	acgtgcagct	780
ttgttacata	ggtaaacgtg	tgccatggtg	gtttgctgta	cctatcaacc	catcacctag	840
gtattaagcc	cagcatgcat	tagctgtttt	tcctgacgct	ctccctctcc	ctgactccca	900
caacaggccc	cagtgtgtgt	tgttcccctc	cctgtgtcca	tgtgttctca	ttgttcagct	960
cccacttata	agtgagaaca	tgtggtgttt	ggttttctgt	ttctgtgtta	gtttgctgag	1020
gataatggct	tccacctcca	tccatgttcc	tgcaaaggac	gtgatcttat	tctttttat	1080
ggttgcatag	aaattgtttt	tacaaatcca	attgatattg	tatttaatta	caagttaatc	1140
taattagcat	actagaagag	attacagaag	atattaggta	cattgaatga	ggaaatatat	1200
aaaataggac	gaaggtgaaa	tattaggtag	gaaaagtata	atagttgaaa	gaagtaaaaa	1260
aaaatatgca	tgagtagcag	aatgtaaaag	aggtgaagaa	cgtaatagtg	actttttaga	1320
ccagattgaa	ggacagagac	agaaaaattt	taaggaattg	ctaaaccatg	tgagtgttag	1380
aagtacagtc	aataacatta	aagcctcagg	aggagaaaag	aataggaaag	gaggaaatat	1440
gtgaataaat	agtagagaca	tgtttgatgg	attttaaaat	atttgaaaga	cctcacatca	1500
aaggattcat	accgtgccat	tgaagaggaa	gatggaaaag	ccaagaagcc	agatgaaagt	1560
tagaaatatt	attggcaaag	cttaaatgtt	aaaagtccta	gagagaaagg	atggcagaaa	1620
tattggcggg	aaagaatgca	gaacctagaa	tataaattca	tcccaacagt	ttggtagtgt	1680

## FIGURE 6B

gcagctgtag	ccttttctag	ataatacact	attgtcatac	atcgcttaag	cgagtgtaaa	1740
atggtctcct	cactttattt	atttatatat	ttatttagtt	ttgagatgga	gcctcgctct	1800
gtctcctagg	ctggagtgca	atagtgcgat	accactcact	gcaacctctg	cctcctctgt	1860
tcaagtgatt	ttcttacctc	agcctcccga	gtagctggga	ttacaggtgc	gtgccaccac	1920
acccggctaa	tttttgtatt	ttttgtagag	acggggtttt	gccatgttgg	ccaggctggt	1980
cttgaactcc	tgacatcagg	tgatccacct	gccttggcct	cctaaagtgc	tgggattaca	2040
ggcatgagcc	accgtgccca	accactttat	ttattttta	tttttattt	taaatttcag	2100
cttctatttg	aaatacaggg	ggcacatata	taggattgtt	acatgggtat	attgaactca	2160
ggtagtgatc	atactaccca	acaggtaggt	tttcaaccca	ctcccctct	tttcctcccc	2220
attctagtag	tgtgcagtgt	ctattgttct	catgtttatg	tctatgtgtg	ctccaggttt	2280
agctcccacc	tgtaagtgag	aacgtgtggt	atttgatttt	ctgtccctgt	gttaattcac	2340
ttaggattat	ggcttccagc	tccattcata	ttgctgtaaa	ggatatgatt	catttttcat	2400
ggccatgcag	tattccatat	tgcgtataga	tcacattttc	tttcttttt	ttttttgaga	2460
cggagtcttg	ctttgctgcc	taggctggag	tgcagtagca	cgatctcggc	tcactgcaag	2520
cttcacctcc	ggggttcacg	tcattcttct	gtctcagctt	cccaagtagc	tgggactaca	2580
ggcgcccgcc	accacgtccg	gctaatttt	ttgtgtgttt	ttagtagaga	tgggggtttc	2640
actgtgttag	ccaggatggt	cttgatctcc	tgaccttgtg	gtccacctgc	ctcggtctcc	2700
caaagtgctg	ggattacagg	ggtgagccac	tgcgcccggc	ccatatatac	cacattttct	2760
ttaaccaatc	caccattgat	gggcaactag	gtagattcca	tggattccac	agttttgcta	2820
ttgtgtgcag	tgtggcagta	gacatatgaa	tgaatgtgtc	tttttggtat	aatgatttgc	2880
attcctttgg	gtatacagtc	attaatagga	gtgctgggtt	gaacggtggc	tctgtttaaa	2940
attctttgag	aattttccaa	actgtttgcc	atagagagca	aactaattta	catttccacg	3000
aacagtatat	aagcattccc	ttttctccac	agctttgtca	tcatggtttt	tttttttctt	3060
tattttaaaa	aagaatatgt	tgttgttttc	ccagggtaca	tgtgcaggat	gtgcaggttt	3120
gttacatagg	tagtaaacgt	gagccatggt	ggtttgctgc	acctgtcaac	ccattacctg	3180
ggtatgaagc	cctgcctgca	ttagctcttt	tccctaatgc	tctcactact	gccccaccct	3240
caccctgaca	gggcaaacag	acaacctaca	gaatgggagg	aaatttttgc	aatctattca	3300
tctgacaaag	gtcaagaata	tccagaatct	acaaggaact	taagcaaatt	tttacttttt	3360

#### FIGURE 6C

aataatagcc actctgactg gcgtgaaatg gtatctcatt gtggttttca tttgaatttc 3420 tetgatgate agtgacgatg ageatttttt catatttqtt qqetqcttqt acqtettttq 3480 agaagtgtct cttcatgcct tttggccact ttaatgggat tattttttgc tttttagttt 3540 aagttootta tagattotgg atattagact tottattgga tgcatagttt gtgaatacto 3600 tettecatte tgtaggttgt etgtttacte tattgatgge ttettttget gtgeegaage 3660 atcttagttt aattagaaac cacctgccaa tttttgtttt tgttgcaatt gcttttgggg 3720 acttagtcat aaactctttg ccaaggtctg ggtcaagaag agtatttcct aggttttctt 3780 ctagaatttt gaaagtctga atgtaaacat ttgcattttt aatgcatctt gagttagttt 3840 tttctttctt tctttctttc tttctttctt tctttctttc tttctttttg tccttctttc 3960 tgcccaggct gcagtgcagc ggcacgatct cggctcactg caacctctqc ctcctqqqtt 4080 caactgattc tectgeatca geetteeaag tagetgggat tataggegee egeeaceaeg 4140 cccgactaat ttttgtattt ttagtagaga cggggttgtg ccatgttggc caggctggtt 4200 tgaaactcct gacctcaaac gatctgcctg ccttggcctc ccaaagtgct gggattacag 4260 gtgtgagcca ctgtgcccag ccaagaatgt cattttctaa gaggtccaag aacctcaaga 4320 tattttggga ccttgagaag agaggaattc atacaggtat tacaagcaca gcctaatggc 4380 aaatctttgg catggcttgg cttcaagact ttaggctctt aaaagtcgaa tccaaaaatt 4440 tttataaaag ctccagctaa gctaccttaa aaggggcctg tatggctgat cactcttctt 4500 gctatacttt acacaaataa acaggccaaa tataatgagg ccaaaattta ttttgcaaat 4560 aaattggtcc tgctatgatt tactcttggt aagaacaggg aaaatagaga aaaatttaga 4620 ttgcatctga ccttttttc tgaattttta tatgtgccta caatttgagc taaatcctga 4680 attattttct ggttgcaaaa actctctaaa gaagaacttg gttttcattg tcttcgtgac 4740 acatttatet ggetetttae tagaacaget ttettgtttt tggtgtteta gettgtgtge 4800 cttacagttc tactcttcaa attattgtta tgtgtatctc atagttttcc ttcttttgag 4860 aaaactgaag ccatggtatt ctgaggacta gagatgactc aacaqagctg gtgaatctcc 4920 tcatatgcaa tccactgggc tcgatctgct tcaaattgct gatgcactgc tgctaaaqct 4980 atacatttaa aacceteact aaaggateag ggaceateat ggaagaggag gaaacatgaa 5040 attgtaagag ccagattcgg ggggtagagt gtggaggtca gagcaactcc accttgaata 5100

## FIGURE 6D

agaaggtaaa	gcaacctatc	ctgaaagcta	acctgccatg	gtggcttctg	attaacctct	5160
gttctaggaa	gactgacagt	ttgggtctgt	gtcattgccc	aaatctcatg	ttaaattgta	5220
atccccagtg	ttcggaggtg	ggacttggtg	gtaggtgatt	cggtcatggg	agtagatttt	5280
cttctttgtg	gtgttacagt	gatagtgagt	gagttctcgt	gagatctggt	catttaaaag	5340
tgtgtggccc	ctcccctccc	tctcttggtc	ctcctactgc	catgtaagat	acctgctcct	5400
gctttgcctt	ctaccataag	taaaagcccc	ctgaggcctc	cccagaagca	gatgccacca	5460
tgcttcctgt	acagcctgca	gaaccatcag	ccaattaaac	ctcttttctg	tataaattac	5520
cagtcttgag	tatctcttta	cagcagtgtg	agaacggact	aatacaaggg	tctccaaaat	5580
tccaagttta	tgtattcttt	cttgccaaat	agcaggtatt	taccataaat	cctgtcctta	5640
ggtcaaacaa	ccttgatggc	atcgtacttc	aattgtctta	cacattcctt	ctgaatgact	5700
cctcccctat	ggcatataag	ccctgggtct	tgggggataa	tggcagaggg	gtccaccatc	5760
ttgtctggct	gccacctgag	acacggacat	ggcttctgtt	ggtaagtctc	tattaaatgt	5820
ttctttctaa	gaaactggat	ttgtcagctt	gtttctttgg	cctctcagct	tcctcagact	5880
ttggggtagg	ttgcacaacc	ctgcccacca	cgaaacaaat	gtttaatatg	ataaatatgg	5940
atagatataa	tccacataaa	taaaagctct	tggagggccc	tcaataattg	ttaagagtgt	6000
aaatgtgtcc	aaagatggaa	aatgtttgag	aactactgtc	ccagagattt	tcctgagttc	6060
tagagtgtgg	gaatatagaa	cctggagctt	ggcttcttca	gcctagaatc	aggagtatgg	6120
ggctgaagtc	tgaagcttgg	cttcagcagt	ttggggttgg	cttccggagc	acatatttga	6180
catgttgcga	ctgtgatttg	gggtttggta	tttgctctga	atcctaatgt	ctgtccttga	6240
ggcatctaga	atctgaaatc	tgtggtcaga	attctattat	cttgagtagg	acatctccag	6300
tcctggttct	gccttctagg	gctggagtct	gtagtcagtg	acccggtctg	gcatttcaac	6360
ttcatataca	gtgggctatc	ttttggtcca	tgtttcaacc	aaacaaccga	ataaaccatt	6420
agaacctttc	cccacttccc	tagctgcaat	gttaaaccta	ggatttctgt	ttaataggtt	6480
catatgaata	atttcagcct	gatccaactt	tacattcctt	ctaccgttat	tctacaccca	6540
ccttaaaaat	gcattcccaa	tatattccct	ggattctacc	tatatatggt	aatcctggct	6600
ttgccagttt	ctagtgcatt	aacatacctg	atttacattc	ttttacttta	aagtggaaat	6660
aagagtccct	ctgcagagtt	caggagttct	caagatggcc	cttacttctg	acatcaattg	6720
agatttcaag	ggagtcgcca	agatcatcct	caggttcagt	gattgctggt	agccctcata	6780

## FIGURE 6E

taactcaatg	aaagctgtta	tgctcatggc	tatggtttat	tacagcaaaa	gaatagagat	6840
gaaaatctag	caagggaaga	gttgcatggg	gcaaagacaa	ggagagctcc	aagtgcagag	6900
attcctgttg	ttttctccca	gtggtgtcat	ggaaagcagt	atcttctcca	tacaatgatg	6960
tgtgataata	ttcagtgtat	tgccaatcag	ggaactcaac	tgagccttga	ttatattgga	7020
gcttggttgc	acagacatgt	cgaccacctt	catggctgaa	ctttagtact	tagcccctcc	7080
agacgtctac	agctgatagg	ctgtaaccca	acattgtcac	cataaatcac	attgttagac	7140
tatccagtgt	ggcccaagct	cccgtgtaaa	cacaggcact	ctaaacaggc	aggatatttc	7200
aaaagcttag	agatgacctc	ccaggagctg	aatgcaaaga	cctggcctct	ttgggcaagg	7260
agaatccttt	accgcacact	ctccttcaca	gggttattgt	gaggatcaaa	tgtggtcatg	7320
tgtgtgagac	accagcacat	gtctggctgt	ggagagtgac	ttctatgtgt	gctaacattg	7380
ctgagtgcta	agaaagtatt	aggcatggct	ttcagcactc	acagatgctc	atctaatcct	7440
cacaacatgg	ctacagggtg	ggcactacta	gcctcatttg	acagaggaaa	ggactgtgga	7500
taagaagggg	gtgaccaata	ggtcagagtc	attctggatg	caaggggctc	cagaggacca	7560
tgattagaca	ttgtctgcag	agaaattatg	gctggatgtc	tctgccccgg	aaagggggat	7620
gcactttcct	tgacccccta	tctcagatct	tgactttgag	gttatctcag	acttcctcta	7680
tgataccagg	agcccatcat	aatctctctg	tgtcctctcc	ccttcctcag	tcttactgcc	7740
cactettece	agctccatct	ccagctggcc	aggtgtagcc	acagtaccta	actctttgca	7800
gagaactata	aatgtgtatc	ctacagggga	gaaaaaaaaa	aagaactctg	aaagagctga	7860
cattttaccg	acttgcaaac	acataagcta	acctgccagt	tttgtgctgg	tagaactcat	7920
gagactcctg	ggtcagaggc	aaaagatttt	attacccaca	gctaaggagg	cagcatgaac	7980
tttgtgttca	catttgttca	ctttgccccc	caattcatat	gggatgatca	gagcagttca	8040
ggtggatgga	cacaggggtt	tgtggcaaag	gtgagcaacc	taggcttaga	aatcctcaat	8100
cttataagaa	ggtactagca	aacttgtcca	gtctttgtat	ctgacggaga	tattatcttt	8160
ataattgggt	tgaaagcaga	cctactctgg	aggaacatat	tgtatttatt	gtcctgaaca	8220
gtaaacaaat	ctgctgtaaa	. atagacgtta	actttattat	ctaaggcagt	aagcaaacct	8280
agatctgaag	gcgataccat	. cttgcaaggc	tatctgctgt	acaaatatgo	: ttgaaaagat	8340
ggtccagaaa	agaaaacggt	attattgcct	ttgctcagaa	gacacacaga	aacataagag	8400
aaccatggaa	aattgtctcc	: caacactgtt	cacccagage	cttccactct	tgtctgcagg	8460
acagtcttaa	catcccatca	ı ttagtgtgtc	: taccacatct	ggcttcacco	g tgcctaacca	8520

#### FIGURE 6F

agatttetag gtecagttee ecaceatgtt tggeagtgee ecactgeeaa ececagaata 8580 agggagtgct cagaattccg aggggacatg ggtggggatc agaacttctg ggcttgagtg 8640 cagagggggc ccatactcct tggttccgaa ggaggaagag gctggaggtg aatgtccttg 8700 gaggggagga atgtgggttc tgaactctta aatccccaag ggaggagact ggtaaggtcc 8760 cagetteega ggtaetgaeg tgggaatgge etgagaggte taagaateee gtateetegg 8820 qaaqqaqqqq ctgaaattqt qaqqqqttga gttgcagggg tttgttagct tgagactcct 8880 tggtgggtcc ctgggaagca aggactggaa ccattggctc cagggtttgg tgtgaaggta 8940 atgggatctc ctgattctca aagggtcaga ggactgagag ttgcccatgc tttgatcttt 9000 ccatctactc cttactccac ttgagggtaa tcacctactc ttctagttcc acaagagtgc 9060 gcctgcgcga gtataatctg cacatgtgcc atgtcccgag gcctggggca tcatccactc 9120 atcattcagc atctgcgcta tgcgggcgag gccggcgcca tgacgtcatg tagctgcgac 9180 tatecetgea gegegeetet eeegteaegt eeeaaceatg gagetgtgga egtgegteee 9240 ctggtggatg tggcctgcgt ggtgccaggc cggggcctgg tgtccgataa agatcctaga 9300 accacaggaa accaggactg aaaggtgcta gagaatggcc atatgtcgct gtccatgaaa 9360 tctcaaggac ttctgggtgg agggcacagg agcctgaact tacgggtttg ccccagtcca 9420 ctgtcctccc aagtgagtct cccagatacg aggcactgtg ccagcatcag cttcatctgt 9480 accacatctt gtaacaggga ctacccagga ccctgatgaa caccatggtg tgtgcaggaa 9540 gagggggtga aggcatggac tcctgtgtgg tcagagccca gagggggcca tgacgggtgg 9600 ggaggaggct gtggactggc tcgagaagtg ggatgtggtt gtgtttgatt tcctttggcc 9660 agataaagtg ctggatatag cattgaaaac ggagtatgaa gaccagttag aatggagggt 9720 caggttggag ttgagttaca gatggggtaa aattctgctt cggatgagtt tggggattgg 9780 caatctaaag gtggtttggg atggcatggc tttgggatgg aaataggttt gtttttatgt 9840 tggctgggaa gggtgtgggg attgaattgg ggatgaagta ggtttagttt tggagataga 9900 atacatggag ctggctattg catgcgagga tgtgcattag tttggtttga tctttaaata 9960 aaggaggeta ttagggttgt ettgaattag attaagttgt gttgggttga tgggttgggc 10020 ttgtgggtga tgtggttgga ttgggctgtg ttaaattggt ttgggtcagg ttttggttga 10080 ggttatcatg gggatgagga tatgettggg acatggattc aggtggttct cattcaagct 10140 gaggcaaatt teettteaga eggteattee agggaaegag tggttgtgtg ggggaaatea 10200

#### FIGURE 6G

ggccactggc tgtgaatatc cctctatcct ggtcttgaat tgtgattatc tatgtccatt 10260 ctqtctcctt cactqtactt qqaattqatc tqqtcattca gctggaaatg ggggaagatt 10320 ttgtcaaatt cttgagacac agctgggtct ggatcagcgt aagcetteet tetggtttta 10380 ttgaacagat gaaatcacat ttttttttc aaaatcacag aaatcttata gagttaacag 10440 tggactetta taataagagt taacaccagg actettatte ttgattettt tetgagacae 10500 caaaatgaga tttctcaatg ccaccctaat tctttttttt tttttttt tttttgagac 10560 acagtctggg tcttttgctc tgtcactcag gctggagcgc agtggtgtga tcatagctca 10620 ctgaaccett gacctectgg acttaaggga teeteetget teageeteet gagtagatgg 10680 qqctacaqqt qcttqccacc acacctgqct aattaaattt ttttttttt tttqtagaga 10740 aagggtetea etttgttgee etggetgate ttgaacttet gaetteaagt gattetteag 10800 cettggactc ccaaagcact gggattgctg gcatgagcca ctcaccgtgc ctggcttgca 10860 gcttaatctt ggagtgtata aacctggctc ctgatagcta gacatttcag tgagaaggag 10920 gcattggatt ttgcatgagg acaattctga cctaggaggg caggtcaaca ggaatccccg 10980 ctgtacctgt acgttgtaca ggcatggaga atgaggagtg aggaggccgt accggaaccc 11040 catattgttt agtggacatt ggattttgaa ataataggga acttggtctg ggagagtcat 11100 atttctggat tggacaatat gtggtatcac aaggttttat gatgagggag aaatgtatgt 11160 ggggaaccat tttctgagtg tggaagtgca agaatcagag agtagctgaa tgccaacgct 11220 tctatttcag gaacatggta agttggaggt ccagctctcg ggctcagacg ggtataggga 11280 ccaggaagtc tcacaatccg atcattctga tatttcaggg catattaggt ttggggtgca 11340 aaggaagtac ttgggactta ggcacatgag actttgtatt gaaaatcaat gattggggct 11400 qqccqtqqtq ctcacqcctq taatctcatc actttqqqaq accqaaqtqq gaggatqqct 11460 tgatctcaag agttggacac cagcctaggc aacatggcca gaccctctct ctacaaaaaa 11520 attaaaaatt agetggatgt ggtggtgcat gettgtggte teagetatee tggaggetga 11580 qacaqqaqaa tcqqttqaqt ctqqqaqttc aaqqctacaq qqaqctqcqa tcacqccqct 11640 qcactccaqc ctqqqaaaca gaqtqagact gtctcagaat ttttttaaaa aagaatcagt 11700 qatcatccca acccctqttq ctqttcatcc tgagcctgcc ttctctggct ttgttcccta 11760 gatcacatct ccatgatcca taggccctgc ccaatctgac ctcacaccgt gggaatgcct 11820 ccagactgat ctagtatgtg tggaacagca agtgctggct ctccctcccc ttccacagct 11880 ctgggtgtgg gaggggttg tccagcctcc agcagcatgg ggagggcctt ggtcagcatc 11940

#### FIGURE 6H

taggtgccaa cagggcaagg gcggggtcct ggagaatgaa ggctttatag ggctcctcag 12000 ggaggcccc cagccccaaa ctgcaccacc tggccgtgga caccggt 12047

cgagcggccc	ctcagcttcg	gcgcccagcc	ccgcaaggct	cccggtgacc	actagagggc	60
gggaggagct	cctggccagt	ggtggagagt	ggcaaggaag	gaccctaggg	ttcatcggag	120
cccaggttta	ctcccttaag	tggaaatttc	ttcccccact	cctccttggc	tttctccaag	180
gagggaaccc	aggctgctgg	aaagtccggc	tggggcgggg	actgtgggtt	caggggagaa	240
cggggtgtgg	aacgggacag	ggagcggtta	gaagggtggg	gctattccgg	gaagtggtgg	300
ggggagggag	cccaaaacta	gcacctagtc	cactcattat	ccagccctct	tatttctcgg	360
ccgctctgct	tcagtggacc	cggggagggc	ggggaagtgg	agtgggagac	ctaggggtgg	420
gcttcccgac	cttgctgtac	aggacctcga	cctagctggc	tttgttcccc	atccccacgt	480
tagttgttgc	cctgaggcta	aaactagagc	ccaggggccc	caagttccag	actgcccctc	540
cccctcccc	cggagccagg	gagtggttgg	tgaaaggggg	aggccagctg	gagaacaaac	600
gggtagtcag	ggggttgagc	gattagagcc	cttgtaccct	acccaggaat	ggttggggag	660
gaggaggaag	aggtaggagg	taggggaggg	ggcggggttt	tgtcacctgt	cacctgctcg	720
ctgtgcctag	ggcgggcggg	cggggagtgg	ggggaccggt	ataaagcggt	aggcgcctgt	780
gecegeteca	cctctcaagc	agecagegee	tgcctgaatc	tgttctgccc	cctccccacc	840
catttcacca	ccaccatq					858

aagcttccac	aagtgcattt	agcctctcca	gtattgctga	tgaatccaca	gttcaggttc	60
aatggcgttc	aaaacttgat	caaaaatgac	cagactttat	attcttacac	caacatctat	120
ctgattggag	gaatggataa	tagtcatcat	gtttaaacat	ctaccattcc	agttaagaaa	180
atatgatagc	atcttgttct	tagtcttttt	cttaataggg	acataaagcc	cacaaataaa	240
aatatgcctg	aagaatggga	caggcattgg	gcattgtcca	tgcctagtaa	agtactccaa	300
gaacctattt	gtatactaga	tgacacaatg	tcaatgtctg	tgtacaactg	ccaactggga	360
tgcaagacac	tgcccatgcc	aatcatcctg	aaaagcagct	ataaaaagca	ggaagctact	420
ctgcaccttg	tcagtgaggt	ccagatacct	acag			454

_	_						_		or Th			_	-	sn Th	cc act nr Thr 15	49
_						gcc Ala						-		_		97
						ttg Leu										145
						att Ile 55										193
aga Arg 65	cgc Arg	gcc Ala	aga Arg	ccc Pro	ccc Pro 70	atc Ile	tat Tyr	agg Arg	cct Pro	atc Ile 75	att Ile	gtg Val	ctc Leu	aac Asn	cca Pro 80	241
cac His	aat Asn	gaa Glu	aaa Lys	att Ile 85	cat His	aga Arg	ttg Leu	gac Asp	ggt Gly 90	ctg Leu	aaa Lys	cca Pro	tgt Cys	tct Ser 95	ctt Leu	289
ctt Leu	tta Leu	cag Gln	tat Tyr 100	gat Asp	taa											307